


Sometimes finding the volume of a solid of revolution is **impossible** by the disk or washer method



Since there is a gap b/w the region and the axis of rotation, we would try washer method

We would have to solve for x as a function of y since the axis of rotation is vertical.

Sometimes this is the problem, but we can do it here.


$$x = \sqrt{\sin^{-1} y}$$

Our problem is that the outer radius and the inner radius use the **same curve**.

In order to find the volume of this solid of revolution we need a different technique.

The **Method of Cylindrical Shells** uses the volume of **nested cylinders** to find the volume of a solid of revolution.

To understand the formula, lets first look at one of the cylindrical shells:



There are two cylinders, an outer cylinder and an inner cylinder.

The volume of the “shell” we use is found by taking the volume of the inner cylinder and subtracting it from the volume of the outer cylinder.

$$V_{shell} = V_{outer} - V_{inner}$$

$$V_{shell} = \pi(r_{outer})^2 h - \pi(r_{inner})^2 h$$

$$V_{shell} = \pi h [(r_{outer})^2 - (r_{inner})^2]$$

$$V_{shell} = \pi h [(r_{outer} + r_{inner})(r_{outer} - r_{inner})]$$


$$V_{shell} = 2\pi h \left(\frac{r_{outer} + r_{inner}}{2} \right) (r_{outer} - r_{inner})$$

$$\frac{(r_{outer} + r_{inner})}{2} = r_{average}$$

Let $r = r_{average}$

Let $\Delta r = r_{outer} - r_{inner}$

$$V_{shell} = \underbrace{2\pi r}_{\text{circumference}} \cdot \underbrace{h}_{\text{height}} \cdot \underbrace{\Delta r}_{\text{thickness}}$$



Math 104 – Rimmer
6.3 Volumes by Cylindrical Shells

$$V_{shell} = 2\pi r \cdot h \cdot \Delta r$$

circumference height thickness

$$V_i = 2\pi(\bar{x}_i) \cdot f(\bar{x}_i) \cdot \Delta x$$


now add up the volume
of all the shells

$$V = \sum_{i=1}^n V_i = \sum_{i=1}^n 2\pi(\bar{x}_i) \cdot f(\bar{x}_i) \cdot \Delta x$$

you get a better approx. as the number of shells $\rightarrow \infty$

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi(\bar{x}_i) \cdot f(\bar{x}_i) \cdot \Delta x$$

$$V = \int_a^b 2\pi x f(x) dx$$



Volume of the solid obtained
by rotating the region under
the curve $f(x)$ from $x = a$ to $x = b$
about the y -axis


In general,


$$V = \int_a^b 2\pi (\text{radius}) (\text{height}) dx$$

distance from axis of rotation to a typical rectangle
height of a typical rectangle
for a vertical axis of rotation dy when the axis is horizontal

Math 104 – Rimmer
6.3 Volumes by Cylindrical Shells

	Typical rectangle	Vertical axis of rotation	Horizontal axis of rotation
Disk or Washer	perpendicular to axis of rotation	Integral is dy (use $x = g(y)$)	Integral is dx (use $y = f(x)$)
Cylindrical Shells	parallel to axis of rotation	Integral is dx (use $y = f(x)$)	Integral is dy (use $x = g(y)$)


Math 104 – Rimmer
6.3 Volumes by Cylindrical Shells



radius = x

height = $\sin(x^2)$

$$V = \int_a^b 2\pi(\text{radius})(\text{height}) dx$$

$$V = 2\pi \int_0^{\sqrt{\pi}} x \sin(x^2) dx$$


$$= 2\pi \left[-\frac{1}{2} \cos x^2 \right]_0^{\sqrt{\pi}}$$

$$= -\pi [\cos \pi - \cos 0]$$

$$= -\pi [-1 - 1]$$

$$= \boxed{2\pi}$$


$u = x^2$
 $du = 2x dx \quad \frac{1}{2} du = x dx$
 $\int \frac{1}{2} \sin u du = -\frac{1}{2} \cos u$


Math 104 – Rimmer
6.3 Volumes by Cylindrical Shells

Set up, but do not evaluate, an integral for the volume obtained by rotating the region bounded by

$y = \cos^2 x$, $y = \frac{1}{4}$, about the line $x = \frac{\pi}{2}$

(below $y = \cos^2 x$ and above $y = \frac{1}{4}$, from $-a$ to a where these are the intersection pts. closest to the y-axis)



radius = $\frac{\pi}{2} - x$

height = $\cos^2 x - \frac{1}{4}$

limits of integration $\Rightarrow x = \frac{\pi}{3} \quad x = -\frac{\pi}{3}$

$\cos^2 x = \frac{1}{4} \Rightarrow \cos x = \frac{1}{2}$

$$V = \int_a^b 2\pi(\text{radius})(\text{height}) dx$$

$$V = \int_{-\pi/3}^{\pi/3} 2\pi \left(\frac{\pi}{2} - x \right) \left(\cos^2 x - \frac{1}{4} \right) dx$$

Website with volumes by shells animation:

<http://mathdemos.gcsu.edu/mathdemos/shellmethod/gallery/gallery.html>

Rihanna rides into the 2008 VMA Awards Show on a shell volume float

Type: “rihanna disturbia 2008 VMA performance”
into the YouTube search, use the first link