



# Math 241 Final Exam Fall 2007

1. Find the radius of convergence for the Taylor series of  $f(z) = \frac{1}{z^8 - 1}$

about the point  $z = 2\sqrt{2} + 2\sqrt{2}i$ .

- (A) 1                      (C) 2                      (E) 3                      (G) 4  
(B)  $\frac{3}{2}$                       (D)  $\frac{5}{2}$                       (F)  $\frac{7}{2}$                       (H)  $\infty$
- 

2. Consider the Laurent series for the function

$$f(z) = \frac{z^2 - 2z + 3}{z - 2}$$

in the region  $|z - 1| > 1$ . What is the coefficient of the  $(z - 1)^{-2}$  term?

- (A) -6                      (C) -3                      (E) 1                      (G) 3  
(B) -4                      (D) 0                      (F) 2                      (H) 6
- 

3. Find the constant  $k$  such that the function  $v(x, y) = 3x^2y + ky^3 - x + 1$  is a harmonic conjugate of the function  $u(x, y) = x^3 - 3xy^2 + y$ .

- (A) -3                      (C) -1                      (E) 1                      (G) 3  
(B) -2                      (D) 0                      (F) 2                      (H) 4
- 

4. Evaluate

$$\int_0^{2i} e^{iz} dz.$$

- (A)  $i(1 - e^{-2})$                       (C)  $1 - ie^{-2}$                       (E)  $i - e^{-2}$                       (G)  $1 - e^{-2}$   
(B)  $i(1 + e^{-2})$                       (D)  $1 + ie^{-2}$                       (F)  $i + e^{-2}$                       (H)  $1 + e^{-2}$
-

5. Evaluate

$$\frac{i}{2} \oint_{|z|=1} \frac{(z^2-1)^2}{z^2(z+\frac{1}{2})(z+2)} dz$$

(A)  $\frac{\pi}{10}$

(C)  $\frac{\pi}{6}$

(E)  $\frac{\pi}{3}$

(G) 1

(B)  $\frac{\pi}{8}$

(D)  $\frac{\pi}{4}$

(F)  $\frac{\pi}{2}$

(H)  $\pi$

---

6. Evaluate

$$\int_0^{2\pi} \frac{d\theta}{2-\cos\theta}$$

(A)  $\pi$

(C)  $\frac{2\pi}{3}$

(E)  $\frac{\pi}{\sqrt{3}}$

(G)  $\frac{\pi}{4}$

(B)  $\frac{2\pi}{\sqrt{3}}$

(D)  $\frac{\pi}{2}$

(F)  $\frac{\pi}{3}$

(H)  $\frac{\pi}{6}$

---

7. Evaluate

$$\int_{-\infty}^{\infty} \frac{dx}{x^2-6x+13}$$

(A)  $-\pi$

(C)  $-\frac{\pi}{24}$

(E)  $\frac{\pi}{24}$

(G)  $\frac{\pi}{2}$

(B)  $-\frac{\pi}{12}$

(D) 0

(F)  $\frac{\pi}{12}$

(H)  $\pi$

---

8. In the Fourier series expansion of  $f(x) = 2x^2 - 1$  on  $(-1,1)$  find the coefficient on the  $\cos(4\pi x)$  term.

- (A) 0                      (C)  $\frac{1}{2\pi}$                       (E)  $\frac{1}{2}$                       (G) 1  
 (B)  $\frac{1}{2\pi^2}$                       (D)  $\frac{2}{\pi^2}$                       (F)  $\frac{2}{\pi}$                       (H) 2
- 

9. Consider the Sturm-Liouville problem defined on  $0 \leq x \leq \frac{\pi}{2}$ :

$$y'' + \lambda y = 0 \quad y(0) = 0, \quad y'\left(\frac{\pi}{2}\right) = 0.$$

Find all eigenvalues  $\lambda_n$ ,  $n = 0, 1, 2, \dots$ .

- (A)  $\lambda_n = n^2$                       (C)  $\lambda_n = \frac{n}{4}$                       (E)  $\lambda_n = \frac{(2n-1)\pi}{4}$                       (G)  $\lambda_n = \frac{(2n-1)^2 \pi}{2}$   
 (B)  $\lambda_n = \frac{n^2}{4}$                       (D)  $\lambda_n = \frac{(2n-1)\pi}{2}$                       (F)  $\lambda_n = (2n-1)^2$                       (H)  $\lambda_n = \frac{(2n-1)^2 \pi}{4}$
- 

10. The solution  $u(x,t)$  defined for  $0 \leq x \leq 2, t \geq 0$  to the wave equation

$$u_{tt} = u_{xx} \quad \text{with boundary conditions } u_x(0,t) = u_x(2,t) = 0 \text{ is}$$

$$u(x,t) = \sum_{n=0}^{\infty} \left[ A_n \cos\left(\frac{n\pi}{2}t\right) + B_n \sin\left(\frac{n\pi}{2}t\right) \right] \cos\left(\frac{n\pi}{2}x\right).$$

Find  $u\left(\frac{1}{3}, \frac{1}{2}\right)$  with initial conditions  $u(x,0) = 3\cos(\pi x)$  and  $u_t(x,0) = 2\cos(3\pi x)$ .

- (A)  $\frac{1}{\pi}$                       (C)  $\frac{3}{\pi}$                       (E)  $\frac{2}{3\pi}$                       (G)  $\frac{1}{2}$   
 (B)  $\frac{2}{\pi}$                       (D)  $\frac{1}{3\pi}$                       (F)  $\frac{1}{3}$                       (H) 2
-

11. Let  $u(x,t)$  be a function defined for  $0 \leq x \leq \pi, t \geq 0$  such that

$$u_t = u_{xx} + 2u_x$$

with boundary conditions  $u(0,t) = u(\pi,t) = 0$  for all  $t \geq 0$

and initial condition  $u(x,0) = e^{-x} \sin(2x)$ .

Use separation of variables to find  $u(\frac{\pi}{4}, 1)$ .

(With separation constant  $-\lambda$ , you will find non-trivial solutions when  $\lambda > 1$ , say  $\lambda = 1 + \alpha^2$ )

- |                            |                             |                            |                             |
|----------------------------|-----------------------------|----------------------------|-----------------------------|
| (A) $e^{-1-\frac{\pi}{2}}$ | (C) $e^{-5-\frac{\pi}{2}}$  | (E) $e^{-1-\frac{\pi}{4}}$ | (G) $e^{-5-\frac{\pi}{4}}$  |
| (B) $e^{-2-\frac{\pi}{2}}$ | (D) $e^{-10-\frac{\pi}{2}}$ | (F) $e^{-2-\frac{\pi}{4}}$ | (H) $e^{-10-\frac{\pi}{4}}$ |
- 

12. Consider a circular plate of radius 1 whose circular edge is maintained at the temperature

$u(1, \theta) = \theta$ . The steady-state temperature  $u(r, \theta) = A_0 + \sum_{n=1}^{\infty} r^n [A_n \cos(n\theta) + B_n \sin(n\theta)]$

satisfies  $u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0$ . Find the coefficient of the  $\sin(3\theta)$  term.

- |                   |                    |                      |                       |
|-------------------|--------------------|----------------------|-----------------------|
| (A) 0             | (C) $\frac{2}{3}$  | (E) $\frac{\pi}{3}$  | (G) $\frac{2\pi}{3}$  |
| (B) $\frac{1}{3}$ | (D) $\frac{-2}{3}$ | (F) $\frac{-\pi}{3}$ | (H) $\frac{-2\pi}{3}$ |
- 

## SOLUTIONS:

1. E
2. G
3. C
4. A
5. H
6. B
7. G
8. B
9. F
10. E
11. G
12. D