

### Problem Set 7

DUE: Thurs. Nov. 4, 2010. *Late papers will be accepted until 1:00 PM Friday.*

**Note:** We say a function is *smooth* if its derivatives of all orders exist and are continuous.

1. Let  $f : [a, \infty) \rightarrow \mathbb{R}$  be a smooth function whose first derivative is bounded:  $|f'(x)| \leq M$  for all  $x \geq a$ . Prove that it is uniformly continuous on  $[a, \infty)$ .

As immediate examples,  $x^{1/3}$  is uniformly continuous for all  $x \geq 1$  and  $\cos x$  is uniformly continuous for all  $x$ .

2. a) Show that  $\sin x$  is not a polynomial.  
 b) Show that  $\sin x$  is not a rational function, that is, it cannot be the quotient of two polynomials.  
 c) Let  $f(t)$  be periodic with period 1, so  $f(t+1) = f(t)$  for all real  $t$ . If  $f$  is not a constant, show that it cannot be a rational function. that is,  $f$  cannot be the quotient of two polynomials.  
 d) Show that  $e^x$  is not a rational function.
3. a) If a smooth function  $f(x)$  has the property that  $f''(x) \geq 0$  for all  $x$ , show that it is *convex*, that is, at every point the graph of the curve  $y = f(x)$  lies above all its tangent lines.  
 b) Let  $v(x)$  be a smooth real-valued function for  $0 \leq x \leq 1$ . If  $v(0) = v(1) = 0$  and  $v''(x) > 0$  for all  $0 \leq x \leq 1$ , show that  $v(x) \leq 0$  for all  $0 \leq x \leq 1$ .  
 c) Prove that the function  $e^x$  is convex.  
 d) Show that  $e^x \geq 1 + x$  for all real  $x$ .
4. a) Let  $p(x) := x^3 + cx + d$ , where  $c$ , and  $d$  are real. Under what conditions on  $c$  and  $d$  does this have three distinct real roots? [ANSWER:  $c < 0$  and  $d^2 < -4c^3/27$ ].  
 b) Generalize to the real polynomial  $p(x) := ax^3 + bx^2 + cx + d$  ( $a \neq 0$ ) by a change of variable  $t = x - \alpha$  (with a clever choice of  $\alpha$ ) to reduce to the above special case.

5. Let

$$p_n(x) := \left(\frac{d}{dx}\right)^n (1-x^2)^n.$$

This is a polynomial of degree  $n$ . Show that it has  $n$  real distinct zeroes, all in the interval  $-1 < x < 1$ .

6. a) For any integer  $n \geq 0$ , show that  $\lim_{x \searrow 0} \frac{e^{-1/x}}{x^n} = 0$ .

b) Define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(x) = \begin{cases} e^{-\frac{1}{x}} & \text{for } x > 0, \\ 0 & \text{for } x \leq 0, \end{cases}$$

Sketch the graph of  $f$ .

c) Show that  $f$  is a smooth function for all real  $x$ .

d) Show that each of the following are smooth and sketch their graphs:

$$g(x) = f(x)f(1-x)$$

$$h(x) = \frac{f(x)}{f(x) + f(1-x)}$$

$$k(x) = h(x)h(4-x)$$

$$K(x) = k(x+2),$$

$$\varphi(x, y) = K(x)K(y), (x, y) \in \mathbb{R}^2$$

$$\Phi(x) = K(\|x\|), x = (x_1, x_2) \in \mathbb{R}^2$$

#### Bonus Problem (Due Nov 4)

B-1 [Interpolation] Let  $x_0 < x_1 < x_2$  be distinct real numbers and  $f(x)$  a smooth function.

a) Show there is a unique quadratic polynomial  $p(x)$  with the property that  $p(x_j) = f(x_j)$  for  $j = 0, 1, 2$ .

b) [Remainder term in interpolation] If  $b$  is in the open interval  $(x_0, x_2)$  with  $b \neq x_j$ ,  $j = 0, 1, 2$ , show there is a point  $c$  (depending on  $b$ ) in the interval  $(x_0, x_2)$  so that

$$f(b) = p(b) + \frac{f'''(c)}{3!}(b-x_0)(b-x_1)(b-x_2).$$

[SUGGESTION: Define the constant  $M$  by

$$f(b) = p(b) + M(b-x_0)(b-x_1)(b-x_2),$$

and look at

$$g(x) := f(x) - [p(x) + M(x-x_0)(x-x_1)(x-x_2)].$$

B-2 In  $\ell_1$  let  $S$  be the set of unit vectors  $e_1 = (1, 0, 0, \dots)$ ,  $e_2 = (0, 1, 0, 0, \dots)$  etc. This set  $S$  is both closed and bounded. Give an example of a continuous  $f : S \rightarrow \mathbb{R}$  that is bounded from above, say  $\sup_{x \in S} f(x) = c$  but there is no point  $p \in S$  where  $f(p) = c$ .

MORAL: In a normed linear space the condition of a set being compact is much stronger than just closed and bounded.

[Last revised: October 28, 2010]