

HOMEWORK 9

1. Prove that $\mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$ is not isomorphic to $\mathbb{Z}/16\mathbb{Z}$ as \mathbb{Z} -modules.
2. How many Abelian groups of size 180 are there up to isomorphism?
3. Consider the ring of $n \times n$ matrices over \mathbb{C} : $R = M_n(\mathbb{C})$. Consider $M_n(\mathbb{C})$ as a module over itself where the addition is the usual addition and scalar multiplication is the usual scalar multiplication. Show that $M_n(\mathbb{C})$ is isomorphic to the direct sum of n copies of \mathbb{C}^n as modules over $M_n(\mathbb{C})$ (where the module structure on \mathbb{C}^n is given by matrix multiplication on column vectors):

$$M_n(\mathbb{C}) \cong \mathbb{C}^n \oplus \dots \oplus \mathbb{C}^n.$$

4. Consider \mathbb{Z} and the multiplicative system $S = \{1, 6, 36, 216, \dots\}$. What is $S^{-1}(\mathbb{Z}/20\mathbb{Z})$ as an $S^{-1}\mathbb{Z}$ -module?
5. Consider \mathbb{Z} and the multiplicative system $S = \mathbb{Z} \setminus 5\mathbb{Z}$. What is $S^{-1}(\mathbb{Z}/20\mathbb{Z})$ as an $S^{-1}\mathbb{Z}$ -module?
6. Consider the group algebra $\mathbb{C}[\mathbb{Z}/3\mathbb{Z}]$ as a left-module over itself. Show that it is isomorphic to the direct sum of three (one-dimensional) submodules $\mathbb{C}_1, \mathbb{C}_2$ and \mathbb{C}_3 where \mathbb{C}_i is generated by the element $1 \cdot \bar{0} + \zeta_3^i \bar{1} + \zeta_3^{2i} \bar{2}$ where ζ_3 is a primitive third root of unity.
7. Find all similarity classes of 3×3 -matrices A over \mathbb{F}_2 so that $A^2 = id$.
8. Find all similarity classes of 3×3 -matrices A over \mathbb{F}_4 so that $A^4 = A$.
9. Let $R_1 = \mathbb{R}[x]$ and V a two-dimensional vector space over \mathbb{R} and fix a basis. Consider the R_1 -module structure on V where x acts as the matrix $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ (in the fixed basis). Prove that V cannot be decomposed as an R_1 -module as a direct sum of 1-dimensional modules.
10. Let $R_2 = \mathbb{C}[x]$ and V a two-dimensional vector space over \mathbb{C} and fix a basis. Consider the R_2 -module structure on V where x acts as the matrix $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ (in the fixed basis). Prove that V can be decomposed as an R_2 -module as a direct sum of 1-dimensional modules.