

# Math 114-003 Fall 2013 Midterm 1

## Exam B

October 29, 2013

Listed below are some practice T/F questions to test your understanding of the material. Please note that actual T/F questions, while similar in style, will not necessarily be variants of the problems below.

1. **True** **False** Consider a differentiable function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that

$$f_x > 0, f_y < 0$$

everywhere. On each level set of  $f$ , the  $y$  coordinate must increase as the  $x$  coordinate increases.

2. **True** **False** For a differentiable function of the form

$$f : \mathbb{R}^3 \rightarrow \mathbb{R},$$

any point of  $f$  neither a local maximum nor a local minimum must be a saddle point.

3. **True** **False** A critical point  $(x_0, y_0)$  of a differentiable function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfying

$$f_{xx}(x_0, y_0) = f_{yy}(x_0, y_0) = 2f_{xy}(x_0, y_0)$$

is necessarily a local minimum.

4. **True** **False** Fix a differential function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $(x_0, y_0) \in \mathbb{R}^2$ . There exists  $C \in \mathbb{R}$  such that

$$|f(x, y) - L(x, y)| < C$$

for all  $x, y$  satisfying  $(x - x_0)^2 + (y - y_0)^2 \leq 7$ , where  $L$  is the standard linear approximation of  $f$  at  $(x_0, y_0)$ .

5. **True** **False** A critical point  $(x_0, y_0)$  of a differentiable function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfying

$$f_{xx}(x_0, y_0), f_{yy}(x_0, y_0) < 0$$

is necessarily a local maximum.

6. **True** **False** The triple integral  $\int \int \int_D x^2 e^y z^4 dV$ , where

$$D = \{(x, y, z) \mid z = x^2 + y^2\},$$

is positive.

7. **True** **False** The double integral  $\int \int_D 16x^2 e^y dA$ , where

$$D = \{(x, y) \mid x^2 + y^2 = 2\},$$

is positive.

8. **True** **False** For a differentiable function  $f : \{(x, y, z) \in \mathbb{R}^3 \mid y > 0\} \rightarrow \mathbb{R}$  defined by

$$f(x, y, z) = e^x - \ln y + x^2 y z^2$$

a unit vector  $\mathbf{u}$  maximizing  $D_{\mathbf{u}}f$  at  $(x_0, y_0, z_0) \in \mathbb{R}^3$  with  $y > 0$  is necessarily

$$\frac{1}{|(\nabla f)(x_0, y_0, z_0)|} (\nabla f)(x_0, y_0, z_0).$$

9. **True** **False** For differentiable functions  $f, g : \mathbb{R}^3 \rightarrow \mathbb{R}$ ,

$$\nabla fg = g\nabla f + f\nabla g.$$

10. **True** **False** For a function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  with  $f_x, f_y, f_{xy}, f_{yx}$  defined and continuous everywhere,

$$f_{xy} = f_{yx}$$

is necessarily true.

11. **True** **False** For a differential function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ , the difference

$$|f(x_0, y_0) - L(x_0, y_0)|,$$

where  $L$  is the standard linear approximation of  $f$  at  $(x_0, y_0)$ , is not necessarily 0.

12. **True** **False** A continuous function  $f : D \rightarrow \mathbb{R}$  must have an absolute minimum if

$$D = \{(x, y, z) \mid |x| + |y| + |z| \leq 5\}.$$