

# Counting Permutations by Their Runs Up and Down

E. Rodney Canfield \*  
University of Georgia  
Athens, GA 30602  
erc@cs.uga.edu

Herbert S. Wilf  
University of Pennsylvania  
Philadelphia, PA 19104-6395  
wilf@math.upenn.edu

October 11, 2006

## Abstract

We find a formula for the number of permutations of  $[n]$  that have exactly  $s$  runs up and down. The formula is at once terminating, asymptotic, and exact.

MSC: 05A15, 05A16

---

\*Research supported by the NSA Mathematical Sciences Program

# 1 Introduction

We will say that a *run* of a permutation  $\sigma$  is a maximal interval of consecutive arguments of  $\sigma$  on which the values of  $\sigma$  are monotonic. If the values of  $\sigma$  increase on the interval then we speak of a run up, else a run down. Throughout this paper we will use the unqualified term *run* to mean either a run up or a run down. These runs have been called *sequences* by some other authors, and have been called *alternating runs* by others. For example, the permutation

$$(723851469)$$

has four runs, viz. 72, 238, 851, 1469. We let  $P(n, s)$  denote the number of permutations of  $n$  letters that have exactly  $s$  runs. Here are the first few values of  $P(n, s)$ :

$n \backslash k$	1	2	3	4
2	2			
3	2	4		
4	2	12	10	
5	2	28	58	32

There is a large literature devoted to this  $P(n, s)$ , which we will survey briefly in section 6. But although a number of recurrences and generating functions etc. are known, it does not seem to have been noticed that an interesting exact formula of the kind we present in this paper exists. Carlitz [6] has derived an exact formula for  $P(n, s)$ , but that one is not at the same time an asymptotic formula. The Carlitz formula is discussed further in the final section.

Our approach to this problem differs from previous studies in that we concentrate on the generating functions  $u_s(x)$ , defined for each fixed  $s \geq 1$  by

$$u_s(x) = \sum_{n \geq 2} P(n, s)x^n,$$

whereas most earlier work has dealt with generating functions for fixed  $n$ . By finding the form of these generating functions we will be able to exhibit a formula for  $P(n, s)$  which is simultaneously

- exact, and
- terminating, and
- asymptotic, for fixed  $s$  and  $n \rightarrow \infty$ .

To our knowledge, the asymptotic behavior of the  $P(n, s)$  has not been previously explored. The formula that we will find is of the form

$$P(n, s) = \frac{s^n}{2^{s-2}} - \frac{(s-1)^n}{2^{s-4}} + \psi_2(n, s)(s-2)^n + \cdots + \psi_{s-1}(n, s), \quad (n \geq 2) \quad (1)$$

in which each  $\psi_i(n, s)$  is a polynomial in  $n$  whose degree in  $n$  is  $\lfloor i/2 \rfloor$ .

## 2 Outline of this paper

In section 3 we will find the generating functions  $u_s(x) = \sum_n P(n, s)x^n$ , as a rational function. Since the denominator will appear in completely factored form, we can write out, in section 4, a formula for  $P(n, s)$  of the type described above.

Interestingly, the formula will be, in that section, uniquely determined except for the coefficient of the leading term! That is, we will show in that section, that for fixed  $s$  we have  $P(n, s) = K(s)s^n + \dots$ , but  $K(s)$  will be, for the moment, unknown.

In section 5 we begin the task of determining the multiplicative factor  $K(s)$ . Surprisingly, although the tools that will have been used up to that point will be entirely analytical in nature, the determination of  $K(s)$  will be done by an “almost-bijection.” We will show that  $P(n, s)$  is, for fixed  $s$ , asymptotic to the number of  $s$ -tuples of pairwise-disjoint subsets of  $[n]$ , each of cardinality  $\geq 2$ , and the asymptotic behavior of the latter is easily found.

The combination of the former analytical results and the latter bijective argument results in the complete formula for  $P(n, s)$ .

## 3 Finding the $u_s(x)$ functions

The recurrence formula for the numbers  $P(n, s)$  is well known and is due to André [1],

$$P(n, s) = sP(n-1, s) + 2P(n-1, s-1) + (n-s)P(n-1, s-2), \quad (n \geq 3) \quad (2)$$

with  $P(2, s) = 2\delta_{s,1}$ . From this recurrence one finds easily a recurrence for the generating functions  $u_s(x) \stackrel{\text{def}}{=} \sum_n P(n, s)x^n$ , viz.

$$(1 - sx)u_s(x) = 2xu_{s-1}(x) + x^2u'_{s-2}(x) - (s-1)xu_{s-2}(x), \quad (s \geq 2) \quad (3)$$

with  $u_1(x) = 2x^2/(1-x)$ ,  $u_0(x) = 0$ . The next three of these functions are

$$\begin{aligned} u_2(x) &= \frac{4x^3}{(1-x)(1-2x)}, \\ u_3(x) &= \frac{2x^4(5-6x)}{(1-3x)(1-2x)(1-x)^2}, \\ u_4(x) &= \frac{4x^5(8-29x+24x^2)}{(1-4x)(1-3x)(1-2x)^2(1-x)^2}. \end{aligned}$$

We will find the general form of these functions, and from that will follow the desired formulas for  $P(n, s)$ .

**Theorem 1** *We have, for each  $s = 1, 2, 3, \dots$ ,*

$$u_s(x) = \frac{\Phi_s(x)}{(1-sx)(1-(s-1)x)(1-(s-2)x)^2(1-(s-3)x)^2 \dots (1-x)^{\lfloor (s+1)/2 \rfloor}}, \quad (4)$$

where  $\Phi_s(x)$  is a polynomial of degree  $1 + \lfloor \frac{s(s+2)}{4} \rfloor$ . The degree of the denominator is  $\lfloor \frac{s(s+2)}{4} \rfloor$ , which is exactly 1 less than the degree of the numerator, for all  $s \geq 1$ .

### 3.1 Proof of Theorem 1

The proof is by a straightforward, though tedious, substitution of the form (4) into the recurrence (3) to find a recurrence for the numerator polynomials  $\Phi_s(x)$ . This will establish that they are indeed polynomials and will provide the claimed degree estimates. We will do this by putting every term over the common denominator

$$\begin{aligned} \Delta_s(x) &= (1-sx)(1-(s-1)x)(1-(s-2)x)^2(1-(s-3)x)^2 \dots (1-x)^{\lfloor (s+1)/2 \rfloor} \\ &\stackrel{\text{def}}{=} \prod_{i=0}^{s-1} (1-(s-i)x)^{\epsilon_i}, \end{aligned}$$

where we have written  $\{\epsilon_i\}_{i \geq 0} = \{1, 1, 2, 2, 3, 3, 4, 4, \dots\}$ .

For technical reasons it will be useful to rewrite the recurrence (3) in the form

$$\begin{aligned}
u_s(x) &= \frac{2xu_{s-1}(x)}{(1-sx)} + \frac{x^2u'_{s-2}(x)}{(1-sx)} - \frac{(s-1)xu_{s-2}(x)}{(1-sx)} \\
&= \frac{2x\Phi_{s-1}(x)}{(1-sx)\Delta_{s-1}(x)} + \frac{x^2\Phi'_{s-2}(x)}{(1-sx)\Delta_{s-2}} - \frac{x^2\Phi_{s-2}}{(1-sx)\Delta_{s-2}} \frac{\Delta'_{s-2}(x)}{\Delta_{s-2}(x)} \\
&\quad - \frac{(s-1)x\Phi_{s-2}(x)}{(1-sx)\Delta_{s-2}(x)} \\
&= \frac{1}{\Delta_s(x)} \left\{ \frac{2x\Phi_{s-1}(x)\Delta_s(x)}{(1-sx)\Delta_{s-1}(x)} + \frac{x^2\Phi'_{s-2}(x)\Delta_s(x)}{(1-sx)\Delta_{s-2}} - \frac{x^2\Phi_{s-2}(x)\Delta_s(x)}{(1-sx)\Delta_{s-2}} \frac{\Delta'_{s-2}(x)}{\Delta_{s-2}(x)} \right. \\
&\quad \left. - \frac{(s-1)x\Phi_{s-2}(x)\Delta_s(x)}{(1-sx)\Delta_{s-2}(x)} \right\} \tag{5}
\end{aligned}$$

Each of the four terms inside the braces is a polynomial in  $x$  whose degree we will now find. Consider the ratio

$$\begin{aligned}
\frac{\Delta_s(x)}{(1-sx)\Delta_{s-1}(x)} &= \frac{\prod_{j=0}^{s-1}(1-(s-j)x)^{\epsilon_j}}{(1-sx)\prod_{j=0}^{s-2}(1-(s-1-j)x)^{\epsilon_j}} \\
&= \frac{\prod_{j=0}^{s-1}(1-(s-j)x)^{\epsilon_j}}{(1-sx)\prod_{j=1}^{s-1}(1-(s-j)x)^{\epsilon_{j-1}}} \\
&= \prod_{j=1}^{s-1} (1-(s-j)x)^{\epsilon_j - \epsilon_{j-1}} = \prod_{j \text{ even}; 2 \leq j \leq s-1} (1-(s-j)x),
\end{aligned}$$

which is a polynomial of degree  $\lfloor (s-1)/2 \rfloor$ .

It follows that

$$\begin{aligned}
\frac{\Delta_s(x)}{(1-sx)\Delta_{s-2}(x)} &= \left( \frac{\Delta_s(x)}{(1-sx)\Delta_{s-1}(x)} \right) \left( \frac{\Delta_{s-1}(x)}{(1-(s-1)x)\Delta_{s-2}(x)} \right) (1-(s-1)x) \\
&= \prod_{j \text{ even}; 2 \leq j \leq s-1} (1-(s-j)x) \prod_{j \text{ even}; 0 \leq j \leq s-2} (1-(s-1-j)x) \\
&= \prod_{j \text{ even}; 2 \leq j \leq s-1} (1-(s-j)x) \prod_{j \text{ odd}; 1 \leq j \leq s-1} (1-(s-j)x) \\
&= \prod_{j=1}^{s-1} (1-(s-j)x),
\end{aligned}$$

is a polynomial in  $x$  of degree  $s-1$ .

We can now deal with the third term of the four inside the braces in (5). Since

$$\frac{\Delta'_{s-2}(x)}{\Delta_{s-2}(x)} = - \sum_{j=2}^{s-1} \frac{\epsilon_{j-2}(s-j)}{1-(s-j)x},$$

we have

$$\frac{x^2 \Phi_{s-2}(x) \Delta_s(x) \Delta'_{s-2}(x)}{(1-sx) \Delta_{s-2}} = x^2 \Phi_{s-2}(x) \left( \prod_{j=1}^{s-1} (1-(s-j)x) \right) \left( \sum_{j=2}^{s-1} \frac{-\epsilon_{j-2}(s-j)}{1-(s-j)x} \right).$$

If  $d(s)$  denotes the degree of  $\Phi(s)$ , then this last member is a polynomial in  $x$  of degree  $2 + d(s-2) + s - 2 = d(s-2) + s$ .

We have now shown that each of the four terms inside the braces in (5) is a polynomial in  $x$ . Their respective degrees are

$$d(s-1) + 1 + \lfloor (s-1)/2 \rfloor, d(s-2) + s, d(s-2) + s, d(s-2) + s.$$

Hence we have  $d(s) = \max(d(s-1) + \lfloor (s+1)/2 \rfloor, d(s-2) + s)$ , with  $d(2) = 3$  and  $d(3) = 5$ .

It is remarkable that this difference equation has a simple solution. Its solution is

$$d(s) = 1 + \left\lceil \frac{s(s+2)}{4} \right\rceil,$$

as can easily be checked, and in fact all four terms inside the braces in (5) have the same degree! This completes the proof of the Theorem.

## 4 The formula for $P(n, s)$

From the partial fraction expansion of (4) we find at once that

$$P(n, s) = \psi_0(n, s)s^n + \psi_1(n, s)(s-1)^n + \psi_2(n, s)(s-2)^n + \cdots + \psi_{s-1}(n, s), \quad (n \geq 2) \quad (6)$$

where each  $\psi_i(n, s)$  is a polynomial in  $n$  of degree at most  $\lfloor i/2 \rfloor$ , and it remains to find these polynomials. We give three methods of doing this: a method of undetermined coefficients, a differential recurrence formula, and finally, a formula of Richard Stanley [11].

## 4.1 Finding the $\psi_i$ 's by undetermined coefficients

Substitute (6) into the recurrence (2) and match the coefficients of each term  $(s-i)^n$ . The result of this substitution is that the  $\psi$ 's satisfy the recurrence

$$(s-i)\psi_i(n, s) = s\psi_i(n-1, s) + 2\psi_{i-1}(n-1, s-1) + (n-s)\psi_{i-2}(n-1, s-2). \quad (7)$$

It should be noted that even if  $\psi_{i-1}$  and  $\psi_{i-2}$  are known, the unknown  $\psi_i$  appears in two places in this recurrence, so we must solve an inhomogeneous difference equation for each  $i$ .

However, we can just assume a solution in the form of a polynomial in  $n$  of degree  $\lfloor i/2 \rfloor$  and solve for the coefficients of that polynomial. We can begin with  $\psi_{-1}(n, s) = 0$  and  $\psi_0(n, s) = K(s)$  (since  $\psi_0$  is of degree zero in  $n$ ) where  $K$  is to be determined. We then find that

$$\psi_1(n, s) = -2K(s-1), \quad \psi_2(n, s) = \frac{1}{4}K(s-2)(s+8-2n)$$

$$\psi_3(n, s) = \frac{1}{2}K(s-3)(2n-s-3), \quad \psi_4(n, s) = \frac{1}{32}K(s-4)(4n^2 - 4n(s+8) + s^2 + 15s + 32)$$

For example, for  $s = 4$  we find

$$u_4(x) = \frac{1/4}{1-4x} + \frac{(-1)}{1-3x} + \frac{(-1/2)}{(1-2x)^2} + \frac{7/2}{1-2x} + \frac{2}{(1-x)^2} + \frac{(-9)}{1-x} + 2x + 19/4.$$

From this it follows

$$P(n, 4) = 4^{n-1} - 3^n + (6-n)2^{n-1} + (2n-7), \quad (n \geq 2).$$

## 4.2 Finding the $\psi_i$ 's recursively

Another method for finding the  $\psi$ 's involves solving the recurrence directly. This leads to a surprisingly elegant differential recurrence, as we will now see. First we need the following lemma about the polynomial solutions of first order inhomogeneous difference equations.

**Lemma 1** *Let  $C \neq 1$ , and let  $f$  be a polynomial. Then the difference equation  $y_{n+1} = Cy_n + f(n)$  has a unique polynomial solution, namely*

$$y_n = -C^{n-1} f\left(x \frac{x}{dx}\right) \left( \frac{x^n}{1-x} \right) \Big|_{x=1/C}. \quad (8)$$

For example, the difference equation  $y_{n+1} = 3y_n + 3n + 2$  has the unique polynomial solution

$$y_n = -3^{n-1} \left( 3x \frac{d}{dx} + 2 \right) \left( \frac{x^n}{1-x} \right) \Big|_{x=1/3} = -\frac{3n}{2} - \frac{7}{4}.$$

To prove the lemma we note first that the general solution of  $y_{n+1} = Cy_n + f(n)$  is evidently

$$y_n = C^n y_0 + \sum_{j=0}^{n-1} C^{n-j-1} f(j) \quad (n = 0, 1, 2, \dots),$$

and we need to discover when this is a polynomial in  $n$ . Now suppose that  $f(n) = \sum_k \alpha_k n^k$ . Then we have

$$y_n = C^n y_0 + C^{n-1} \sum_k \alpha_k \sum_{j=0}^{n-1} C^{-j} j^k.$$

But it is easy to check by induction that

$$\sum_{j=0}^{n-1} j^k x^j = \left( x \frac{d}{dx} \right)^k \left( \frac{1-x^n}{1-x} \right) = x^n \frac{R_k(x, n)}{(x-1)^{k+1}} + \frac{Q_k(x)}{(x-1)^{k+1}}, \quad (9)$$

where  $R_k$  is a polynomial in  $x$  of degree  $k$  and is also a polynomial in  $n$  of degree  $k$ , and  $Q_k$  is a polynomial in  $x$  of degree  $k$ . Consequently the general solution is

$$y_n = C^n y_0 + C^{n-1} \sum_k \alpha_k \frac{Q_k(C^{-1})}{(C^{-1}-1)^{k+1}} + C^{-1} \sum_k \alpha_k \frac{R_k(C^{-1}, n)}{(C^{-1}-1)^{k+1}}.$$

Since  $C \neq 1$ , this will be a polynomial in  $n$  iff

$$y_0 = -C^{-1} \sum_k \alpha_k \frac{Q_k(C^{-1})}{(C^{-1}-1)^{k+1}},$$

and if that condition is satisfied the unique polynomial solution will be

$$y_n = \frac{1}{C} \sum_k \alpha_k \frac{R_k(C^{-1}, n)}{(C^{-1}-1)^{k+1}}.$$

We can simplify the form of this answer by recalling that, from (9) we have

$$\frac{R_k(x, n)}{(x-1)^{k+1}} = -x^{-n} \left( x \frac{d}{dx} \right)^k \left( \frac{x^n}{1-x} \right),$$

and therefore we can cast the unique polynomial solution in the more pleasing form (8).  $\square$

Hence we have the following procedure for calculating the  $\psi$ 's. For each  $i = 1, 2, \dots$  we do

1. Suppose  $\psi_{i-2}$  and  $\psi_{i-1}$  are known.
2. Define the polynomial

$$f(n) = \frac{2}{s-i} \psi_{i-1}(n, s-1) + \frac{(n+1-s)}{s-i} \psi_{i-2}(n, s-2),$$

and put  $C = s/(s-i)$ .

3. Then

$$\psi_i(n, s) = -C^{n-1} f\left(x \frac{x}{dx}\right) \left(\frac{x^n}{1-x}\right) \Big|_{x=1/C}.$$

### 4.3 Finding the $\psi_i$ 's from Stanley's formula

In [11] Richard Stanley has given an exact formula for  $P(n, k)$ , viz.

$$P(n, k) = \sum_{\ell=0}^k (-1)^{k-\ell} \frac{z_{k-\ell}}{2^{\ell-1}} \sum_{\substack{r+2m \leq \ell \\ r \equiv \ell \pmod{2}}} (-2)^m \binom{\ell-m}{(\ell+r)/2} \binom{n}{m} r^n,$$

where  $z_0 = 2$  and all other  $z_i$ 's are 4. Evidently this contains an implicit formula for our  $\psi$ 's.

## 5 The factor $K(s)$

We have now described the formula for  $P(n, s)$  completely except for the multiplicative factor  $K(s)$ . It remains to show that  $K(s) = 2^{-(s-2)}$ . For this, it would suffice to prove the next Theorem for fixed  $s$  and  $n \rightarrow \infty$ ; since the proof is applicable to a larger range of  $s$ , we state it in that manner:

**Theorem 2** *Let  $\epsilon > 0$ , and  $\{(n, s)\}$  be an infinite sequence of pairs such that  $n \rightarrow \infty$  and  $s \leq (1 + \epsilon)^{-1} n / \log n$ . Then,*

$$P(n, s) \sim \frac{1}{2^{s-2}} s^n. \tag{10}$$

### 5.1 Proof of Theorem 2

To fix ideas, we will do this by showing that the number  $\hat{P}(n, s)$  of permutations of  $n$  letters, with  $s$  runs, *the first of which is a run up*, is  $\sim s^n / 2^{s-1}$ . Evidently the number for which the first run is down will be the same, and the desired result will follow. Henceforth we will

always assume that the first run is a run up. There are two steps to the proof. In the first step, we show that the set of permutations counted by  $\hat{P}(n, s)$  can be put into bijection with  $s$ -tuples of subsets  $(S_1, \dots, S_s)$  (each  $S_i \subseteq [n]$ ) satisfying certain properties. In the second part of the proof, we introduce a function called  $\Phi$  whose domain is the Cartesian product of these  $s$ -tuples with a set of cardinality  $2^{s-1}$ , and whose range is a set of size  $s^n$ . We prove that this function  $\Phi$  is an *injection*. Although we have no succinct description of the image of this injection, we are able to show that for  $(n, s)$  in the range hypothesized by the theorem the image is asymptotically all of the range set.

## 5.2 First part of the proof

Let  $\Pi(n, s)$  be the set of all  $n$ -permutations with  $s$  runs up and down, the first of which is up. Let  $\tilde{\Pi}(n, s)$  be the collection of all  $s$ -tuples  $(S_1, \dots, S_s)$  of nonempty subsets of  $[n]$  which are *almost* pairwise disjoint, in that

$$|S_i \cap S_j| = \begin{cases} 1, & \text{if } j = i + 1 \text{ and } 1 \leq i < s; \\ 0, & \text{else} \end{cases} \quad (11)$$

Further we require that

$$|S_i| \geq 2, \quad \forall i, \quad (12)$$

and that

$$\begin{aligned} \max(S_i) &= \max(S_{i+1}) \in S_i \cap S_{i+1} && (\forall \text{ odd } i) \\ \min(S_i) &= \min(S_{i+1}) \in S_i \cap S_{i+1} && (\forall \text{ even } i). \end{aligned} \quad (13)$$

**Lemma 2** *The number of  $s$ -tuples of subsets of  $[n]$  that satisfy (11)–(13) is equal to the number of permutations of  $[n]$  with  $s$  runs, the first of which is up.*

Indeed to reconstruct the permutation from the  $s$ -tuple of sets, we first sort each of the sets, the first in increasing order, the second decreasing, etc., then merge the sets, and finally delete one element of each of the adjacent duplicates that appear.  $\square$

Hence it suffices to show that the number of  $s$ -tuples of subsets that satisfy (11)–(13) is  $\sim s^n / 2^{s-1}$ .

## 5.3 Defining the function $\Phi$

By a *choice sequence*  $\mathbf{h} = (h_1, \dots, h_{s-1})$  we mean an  $s - 1$ -tuple where each  $h_i$  is either equal to  $i$  or to  $i + 1$ . The set of all such choice sequences will be  $H_s$ . The function to be

constructed is a mapping

$$\Phi : H_s \times \tilde{\Pi}(n, s) \rightarrow \{(T_1, T_2, \dots, T_s) : \forall i, T_i \subseteq [n]\}.$$

Let  $\mathbf{h} \in H_s$ , and let  $(S_1, \dots, S_s)$  be a family of subsets satisfying (11)–(13). For each  $i = 1, \dots, s - 1$ , let  $e_i$  be the unique element that belongs to  $S_i \cap S_{i+1}$ . These  $e_i$ 's are all different, since  $e_i = e_j$  with  $i < j$  would imply that  $S_i \cap S_{j+1}$  is nonempty, contradicting (11). Perform the following  $s - 1$  delete operations: for each  $i = 1, \dots, s - 1$ , delete the element  $e_i$  from the set  $S_{h_i}$ . The resulting  $s$ -tuple of sets remaining after these deletions is, by definition,  $\Phi(\mathbf{h}, (S_1, \dots, S_s))$ .

The image of this mapping does not include all  $s$ -tuples of sets, as the following Lemma shows.

**Lemma 3** *If  $(T_1, \dots, T_s)$  is in the image of  $\Phi$  then*

1. *the  $T_i$ 's are pairwise disjoint, and*
2. *the union of the  $T_i$ 's is  $[n]$ .*

□

It is possible for some of the  $T_i$ 's to be empty. We remark that the number of  $s$ -tuples  $(T_1, \dots, T_s)$  in which the  $T_i$ 's are pairwise disjoint and whose union is  $[n]$  is  $s^n$ .

## 5.4 The mapping $\Phi$ is injective

The way we prove this assertion is to give a *reconstruction* algorithm. The algorithm begins with an  $s$ -tuple  $(T_1, \dots, T_s)$  of subsets which putatively belongs to the image of  $\Phi$ . It attempts to reconstruct the preimage. It will be clear from the algorithm that the preimage can be only one thing, if it exists at all. There is one “early exit” point in the algorithm where the search for a preimage is abandoned, because it obviously does not exist. If the algorithm executes all the way to finish, then it will have found the only possible candidate for a preimage. However, it is still possible that the  $s$ -tuple of sets found at the end will not satisfy one of the required conditions (11)–(13).

**Lemma 4** *The mapping  $\Phi$  is injective.*

**Proof.** Let  $(T_1, \dots, T_s)$  be an  $s$ -tuple of pairwise disjoint (possibly empty) sets whose union is  $[n]$ . Here is the reconstruction algorithm:

1. (Find consecutive unions) It is easy to see that if the  $T$ 's are in the image of  $\Phi$ , then

$$T_i \cup T_{i+1} = S_i \cup S_{i+1}, \quad 1 \leq i < s.$$

So if one of the inequalities

$$|T_i \cap T_{i+1}| \geq 3 \quad 1 \leq i < s,$$

fails, then the reconstruction fails and no preimage exists. Otherwise we can reconstruct all of the unions  $S_i \cup S_{i+1}$ .

2. (Reconstruct the set of deleted elements) Put  $e_1 = \max(S_1 \cup S_2)$ ,  $e_2 = \min(S_2 \cup S_3)$ ,  $\dots$
3. (Recover the choice sequence  $\mathbf{h}$ ) For each  $i = 1, \dots, s - 1$ , since  $e_i \in T_i \cup T_{i+1}$ , and because the  $T_i$ 's are pairwise disjoint, there will be exactly one index,  $h_i$ , say, such that  $h_i \in \{i, i + 1\}$  and  $e_i \notin T_{h_i}$ .
4. (Re-insert the elements that were deleted) For each  $i$ ,  $1 \leq i < s$ , insert the element  $e_i$  into the set  $T_{h_i}$ .

If the reconstructed sets  $(S_1, \dots, S_s)$  satisfy (11)–(13) then we have found the unique preimage. Otherwise no preimage exists.  $\square$

Thus if  $\hat{P}(n, s)$  is the number of permutations of  $n$  letters with  $s$  runs, the first of which is up, then we have shown that

$$2^{s-1} \hat{P}(n, s) \leq s^n. \tag{14}$$

## 5.5 When does the algorithm terminate without a preimage?

If the reconstruction algorithm does not early exit in step 1, yet fails to find a preimage, then one of the conditions (11)–(13) is not satisfied. We will now visit each of these in turn to see when it might fail

1. (Can (11) fail?) The intersections  $T_i \cap T_{i+1}$  were all empty before the insertions; however, the operation, “insert element  $e_i$  into  $T_{h_i}$ ” either added an element of  $T_i$  to the set  $T_{i+1}$ , or vice-versa. That operation alone caused the two adjacent sets to have intersection 1. The only other insertion which could have affected  $T_i$  is the one which involves element  $e_{i-1}$ . If that operation increased the size of  $T_i$ , then it did so by inserting an element from  $T_{i-1}$ , which element could not possibly be present in  $T_{i+1}$ . Thus, the only other insertion which could possibly affect the set  $T_i$  will have no effect on the cardinality of  $T_i \cap T_{i+1}$ . Likewise, the only other operation which can possibly affect the cardinality of  $T_{i+1}$  will have no effect on the cardinality of  $T_i \cap T_{i+1}$ . So, the intersection  $S_i \cap S_{i+1}$  will always have size 1, as required.

2. (Will  $S_i \cap S_j = \emptyset$  when  $j > i + 1$ ?) Only the case  $j = i + 2$  is not obvious. If  $S_i \cap S_{i+2}$  is not empty, then during reconstruction some element originally belonging to  $T_{i+1}$  was inserted into both  $T_i$  and  $T_{i+2}$ . (Any element originally in  $T_i$  cannot end up in  $S_{i+2}$ , and vice-versa.) This means that some element  $e \in T_{i+1}$  is both the maximum of  $T_i \cup T_{i+1}$ , as well as the minimum of  $T_{i+1} \cup T_{i+2}$ ; (or the other way around). But

$$\max(T_{i+1}) \leq \max(T_i \cup T_{i+1}) = e_i = \min(T_{i+1} \cup T_{i+2}) \leq \min(T_{i+1}),$$

so  $T_{i+1}$  has just one element,  $e_i$ , and that element lies between the surrounding sets  $T_i$  and  $T_{i+2}$ . This is how the reconstructed sets can fail to satisfy (11).

3. (Can (12) fail?) Yes, if one of the sets  $T_i$  has cardinality 0 or 1, then it is possible that the cardinality of the reconstructed  $S_i$  may be less than 2.
4. (Can (13) fail?) No. By the nature of the reconstruction, the  $S_i$ 's always have this property.

We can now prove

**Lemma 5** *If in the given sequence  $T = (T_1, \dots, T_s)$ , all sets have cardinalities at least 2, then  $T$  has a preimage under  $\Phi$ .*

For then the unions  $T_i \cup T_{i+1}$  have size 4 or more, so we don't terminate the reconstruction at Step 1. The only other two possible failures — when an intersection  $S_i \cap S_{i+2}$  was nonempty, or one of the  $S_i$  was too small — could both be traced back to a set  $T_i$  which had size 0 or 1.  $\square$

A crude lower estimate from Bonferroni's inequalities tells us that the number of  $s$ -tuples  $T$  that are pairwise disjoint, with union equal to  $[n]$ , and with all cardinalities  $\geq 2$  is at least

$$s^n - (n + s)(s - 1)^{n-1}.$$

The reason:  $s(s - 1)^n$  is an upper bound on the number of  $T$ 's for which some component is the empty set; and  $ns(s - 1)^{n-1}$  is an upper bound on the number of  $T$ 's for which some component has cardinality one; then,  $s(s - 1) + ns < s(n + s)$ . Hence

$$2^{s-1} \hat{P}(n, s) \geq s^n - s(n + s)(s - 1)^{n-1}, \quad (15)$$

which, taken together with (14) completes the proof of (10), since our hypothesis on the pairs  $(n, s)$  implies

$$(n + s)(s - 1)^{n-1} \leq 2n(s - 1)^{n-1} = o(s^n).$$

## 6 Survey of the literature

André was the first to study [1] the runs up and down of permutations, and the fundamental recurrence (2) is due to him. His paper includes a table of  $P(n, s)$  through  $n = 8$ , with one error in the final row. A great deal of information about  $P(n, s)$  is found in vol. 3 of [9] (see particularly ex. 15, 16 of sec. 5.1.3).

The history of generating functions in this problem is complex. Comtet [8] devotes an extended exercise, see page 260, to the topic. The two variable generating function given there, however, is incorrect. Carlitz [5, 6, 7] visited this subject several times. In [5] he gives a two-variable generating function

$$\sum_{n=2}^{\infty} \frac{z^n}{n!} (1-x^2)^{-n/2} \sum_{s=1}^{n-1} P(n+1, s) x^{n-s} = \frac{(1-x)((1-x^2)^{1/2} + \sin(z))^2}{(1+x)(x - \cos(z))^2},$$

and in [6] he finds an explicit formula for  $P(n, s)$  and information about an associated polynomial sequence. There is something wrong with the final formulas of this latter work, however; these formulas suggest  $P(8, s) = 0, 2, 250, 2516, 7060, 7562, 2770$ ; whereas, in fact,  $P(8, s) = 2, 252, 2766, 9576, 14622, 10332, 2770$ . (Empirically, his formula always gives the right value for  $P(n, n-1)$ .) Further evidence that something is amiss concerns the auxiliary quantity  $\overline{K}_{n,j}$ ; the summation formula given for this quantity does not give the values displayed in the table. Whether the problem can be easily repaired, we have not investigated.

A correct generating function appears in the discussion accompanying sequence A059427 of [10]. This one is due to Emeric Deutsch and Ira Gessel, who sent it to Sloane in December of 2004. A correct generating function is also in Stanley [11], who used an observation of Miklós Bóna to connect the sequence that we study here with  $a_k(n)$ , the number of  $n$ -permutations the length of whose longest alternating subsequence is  $k$ .

Bóna and Ehrenborg [4] have proven log-concavity:  $P(n, s)^2 \geq P(n, s-1)P(n, s+1)$ . In the later book [3], the stronger assertion, that  $P_n(x) \stackrel{\text{def}}{=} \sum_s P(n, s)x^s$  has all its roots real and negative, is made. A proof of this can be based on the relation

$$P_n(x) = (x - x^3)P'_{n-1}(x) + ((n-2)x^2 + 2x)P_{n-1}(x),$$

which itself is a consequence of the basic recursion (2). This implies, once it is established that the variance becomes infinite with  $n$ , that the numbers  $P(n, s)$  satisfy a central limit theorem. (That is, are asymptotically normal.) Due to log-concavity, one may deduce (see Theorem 4 of [2]) a local limit theorem. This leads to an asymptotic formula for  $P(n, s)$  for  $s$  in a different range than in our Theorem 2.

## References

- [1] M. Désiré André, Étude sur les maxima, minima et séquences des permutations, Ann. Sci. École Norm. Sup. **3** no. 1 (1884), 121–135.
- [2] E. A. Bender, Central and local limit theorems applied to asymptotic enumeration, J. Combinatorial Theory, series A **15** (1973) 91–111.
- [3] M. Bóna, *The Combinatorics of Permutations*, Chapman and Hall/CRC, 2004.
- [4] M. Bóna and R. Ehrenborg, A combinatorial proof of the log-concavity of the numbers of permutations with  $k$  runs. J. Combinatorial Theory, series A **90** (2000) 293–303.
- [5] Carlitz, Leonard, Enumeration of permutations by sequences. Fibonacci Quart. **16** (1978), no. 3, 259–268.
- [6] Carlitz, L. The number of permutations with a given number of sequences. Fibonacci Quart. **18** (1980), no. 4, 347–352.
- [7] Carlitz, L. Enumeration of permutations by sequences. II. Fibonacci Quart. **19** (1981), no. 5, 398–406, 465.
- [8] Louis Comtet, *Advanced Combinatorics*, Reidel, 1974.
- [9] Knuth, Donald E., *The Art of Computer Programming*, vol. 3, Addison Wesley, 1973.
- [10] Sloane, Neil, The On-Line Encyclopedia of Integer Sequences, seq. A059427; on the web at <http://www.research.att.com/~njas/sequences>.
- [11] Stanley, Richard P., Longest alternating subsequences of permutations, arXiv:math.CO/0511419v1 (Nov. 16, 2005).