

Name: _____

Midterm Exam II for Math 110, Fall 2015

November 02, 2017

Problem	Points	Score
1	15	
2	16	
3	15	
4	12	
5	12	
6	18	
7	12	
8	12	
9	8	
Total	120	

- Please show **ALL** your work on this exam paper. Partial credit will be awarded where appropriate.
- **CLEARLY** indicate final answers. Use words (doesn't have to be that many words) to connect mathematical formulas and equations.
- **NO** books, notes, laptops, cell phones, calculators, or any other electronic devices may be used during the exam. One 8.5×11 cheatsheet, handwritten is allowed; it may be double-sided.
- No form of cheating will be tolerated. You are expected to uphold the Code of Academic Integrity.

My signature below certifies that I have complied with the University of Pennsylvania's Code of Academic Integrity in completing this midterm examination.

Signature: _____

Date: _____

1. (a) Find c and p such that $\frac{x^{-2}}{2 + 1/x} \sim cx^p$ as $x \rightarrow \infty$.

$$c = \underline{\hspace{2cm}}$$

$$p = \underline{\hspace{2cm}}$$

Show your work or justify your answer.

- (b) Write this integral as a limit.

$$\int_1^{\infty} \frac{x^{-2}}{2 + 1/x} dx$$

- (c) Say (with justification) whether or not the integral in part (b) is convergent.

2. (a) Compute the indefinite integral of x^2e^{-x} .

(b) Write $\int_0^\infty x^2e^{-x} dx$ as a limit and compute its value using the indefinite integral you computed in part (a). Please briefly justify any assertions about the limit as $M \rightarrow \infty$.

(c) Compute the value of C that makes Cx^2e^{-x} a probability density on the positive real numbers.

(d) Compute the mean of this density.

4. Write a differential equation for this scenario. You do not have to solve the differential equation but you must give the interpretation of all variables and constants, their units, and indicate which is the dependent and the independent variable.

The amount of an endowment increases due to investment at a rate proportional to the value, decreased by fees that are taken out at a constant rate.

5. (a) For which values of x does this series converge?

$$\sum_{n=0}^{\infty} \frac{2^n}{n!} x^n$$

(b) Compute the MacLaurin series for e^{2x} and state its radius of convergence.

6. A revenue forecast for a technological innovation forecasts earnings of $n^{-2/5}$ million dollars in year n .

(a) Express the total possible earnings of this innovation (assuming you can continue marketing it forever) as an infinite series in Sigma-notation.

(b) Say whether this total is finite or infinite.

(c) **If you answered “infinite”:**

Approximately how much should you have earned after some large number of years, M ? Use an integral approximation and write your answer as cM^p .

If you answered “finite”:

Approximately how much value remains to be earned after M years? Use an integral approximation and write your answer as cM^p .

$$c = \underline{\hspace{2cm}}$$

$$p = \underline{\hspace{2cm}}$$

7. (a) Choose which initial value problem corresponds to this integral equation.

$$y(x) = 5x^2 + \int_3^x \frac{1}{y(t)} dt.$$

- (A) $y' = 5x^2 + 1/y$; $y(0) = 0$
- (B) $y' = 5x^2 + 1/y$; $y(3) = 30$
- (C) $y' = 5x^2 + 1/y$; $y(3) = 45$
- (D) $y' = 5x^2 + 1/y$; $y(0) = 5x^2$
- (E) $y' = 10x + 1/y$; $y(0) = 0$
- (F) $y' = 10x + 1/y$; $y(3) = 30$
- (G) $y' = 10x + 1/y$; $y(3) = 45$
- (H) $y' = 5x^2 + y'/y$; $y(3) = 0$

- (b) Choose which integral equation corresponds to this initial value problem.

$$y' = \frac{x}{1+y} ; y(2) = 17.$$

- (A) $y(x) = \int x/(1+y) dx$; $y(2) = 17$
- (B) $y(x) = \int_0^\infty x/(1+y) dx$
- (C) $y(x) = \int_1^x t/(1+y) dt$; $y(2) = 17$
- (D) $y(x) = 17 + \int_2^x t/(1+y) dt$
- (E) $y(x) = 17 + \int_0^x t/(1+y) dt$
- (F) $y(x) = \int_0^x t/(1+y) dt + \int_0^2 17 dx$

8. Use integrals (hint: you may need to integrate by parts) to find upper and lower bounds on this sum:

$$\sum_{k=1}^{100} \ln(k).$$

Express these bounds as exact quantities involving the natural log. Evaluate one of them approximately by using the log cheatsheet.

9. Which of these values comes closest to $y(3)$ if $y(x)$ satisfies the differential equation $y' = (10 + x)\frac{y}{y+1}$ with initial condition $y(1) = 50$? Circle one answer. If you wish to be considered for partial credit, then provide justification.

(A) 0

(B) $72/52$

(C) 50

(D) 70

(E) 90

(F) 5,000

TABLE 8.1 Basic integration formulas

1. $\int k \, dx = kx + C$ (any number k)	12. $\int \tan x \, dx = \ln \sec x + C$
2. $\int x^n \, dx = \frac{x^{n+1}}{n+1} + C$ ($n \neq -1$)	13. $\int \cot x \, dx = \ln \sin x + C$
3. $\int \frac{dx}{x} = \ln x + C$	14. $\int \sec x \, dx = \ln \sec x + \tan x + C$
4. $\int e^x \, dx = e^x + C$	15. $\int \csc x \, dx = -\ln \csc x + \cot x + C$
5. $\int a^x \, dx = \frac{a^x}{\ln a} + C$ ($a > 0, a \neq 1$)	16. $\int \sinh x \, dx = \cosh x + C$
6. $\int \sin x \, dx = -\cos x + C$	17. $\int \cosh x \, dx = \sinh x + C$
7. $\int \cos x \, dx = \sin x + C$	18. $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + C$
8. $\int \sec^2 x \, dx = \tan x + C$	19. $\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$
9. $\int \csc^2 x \, dx = -\cot x + C$	20. $\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1}\left \frac{x}{a}\right + C$
10. $\int \sec x \tan x \, dx = \sec x + C$	21. $\int \frac{dx}{\sqrt{a^2 + x^2}} = \sinh^{-1}\left(\frac{x}{a}\right) + C$ ($a > 0$)
11. $\int \csc x \cot x \, dx = -\csc x + C$	22. $\int \frac{dx}{\sqrt{x^2 - a^2}} = \cosh^{-1}\left(\frac{x}{a}\right) + C$ ($x > a$)

Logarithm Cheat Sheet

These values are accurate to within 1%:

$$e \approx 2.7$$

$$\ln(2) \approx 0.7$$

$$\ln(10) \approx 2.3$$

$$\log_{10}(2) \approx 0.3$$

$$\log_{10}(3) \approx 0.48$$

Some other useful quantities to with 1%:

$$\pi \approx \frac{22}{7}$$

$$\sqrt{10} \approx \pi$$

$$\sqrt{2} \approx 1.4$$

$$\sqrt{1/2} \approx 0.7$$

(ok so technically $\sqrt{2}$ is about 1.005% greater than 1.4 and 0.7 is about 1.005% less than $\sqrt{1/2}$)