MINIMAL HYPERSURFACES WITH BOUNDED INDEX

OTIS CHODOSH, DANIEL KETOVER, AND DAVI MAXIMO

ABSTRACT. We prove a structural theorem that provides a precise local picture of how a sequence of closed embedded minimal hypersurfaces with uniformly bounded index (and volume if the ambient dimension is greater than three) in a Riemannian manifold (M^n, g) , $3 \le n \le 7$, can degenerate. Loosely speaking, our results show that embedded minimal hypersurfaces with bounded index behave qualitatively like embedded stable minimal hypersurfaces, up to controlled errors. Several compactness/finiteness theorems follows our local picture.

1. INTRODUCTION

Minimal hypersurfaces are critical points of the volume functional, and as such it is natural to study their existence and behavior from a variational point of view. A key invariant related to this point of view is the (Morse) index of such an object; and the assumption of bounded index (rather than genus or total curvature as in the classical theory) that we are concerned with in this paper is very natural. Many minimal surfaces with bounded index are expected to arise from the variational min-max theory of Almgren–Pitts [Pit76]. For instance, Marques–Neves [MN13] have introduced nontrivial k-parameter sweep-outs in arbitrary three-manifolds (for any k), and the Morse index of the corresponding minimal surface is expected (generically) to be k. Moreover, Colding–Gabai [CG14] have recently studied sequences of index one minimal surfaces and as they relate to the problem of classifying Heegaard splittings of three-manifolds.

In this work, we provide a precise local picture of how a sequence of embedded minimal hypersurfaces with uniformly bounded index (and volume if the ambient dimension is greater than three) in a Riemannian manifold (M^n, g) , $3 \le n \le 7$, can degenerate. We may roughly describe it as follows. For the sake of exposition, we assume here that n = 3, the surfaces are all two-sided and have uniformly bounded area, *i.e.*, consider a sequence of embedded two-sided minimal surfaces Σ_j in a closed Riemannian (M^3, g) so that $index(\Sigma_j) \le I$ and $area_g(\Sigma_j) \le \Lambda$. Given these assumptions, our results imply that $genus(\Sigma_j)$ is uniformly bounded:

- (1) A blow-up argument allows us to extend Schoen's curvature estimates [Sch83a] to the case of bounded index (cf. Corollary 2.3), to see that the curvature of Σ_j is bounded away from at most I points, where the index may be concentrating. Blowing up around these points at the scale of curvature, we produce a smooth embedded minimal surface in \mathbb{R}^3 with index at most I.
- (2) By [Ros06] (see also [CM14]), this limiting surface has genus bounded linearly above in terms of I. This can be seen as a kind of "lower semi-continuity of

Date: December 8, 2015.

topology," since each genus that is seen in the blow-up limit certainly contributes to the genus of Σ_j for j sufficiently large.

(3) Furthermore, after passing to a subsequence, Σ_j converges to a closed minimal surface in (M, g) smoothly away from the points of index concentration. Such a surface has bounded genus.

These facts, by themselves, are not sufficient to conclude that the genus of Σ_j is uniformly bounded. The reason for this is that it is a priori possible that some genus is lost to the intermediate scales, and thus does not appear in the blow-up limits, or at the original scale. This can be illustrated by an analogy with the bubbling phenomenon for harmonic maps; a sequence of harmonic maps may degenerate to form a bubble tree and the key point in proving that the energy of the bubbles is the limit of the energy of the original sequence of maps, is to show that no energy is lost in the neck regions joining the bubbles. Hence, to prove genus bounds, we must also show:

(4) No topology is lost in the intermediate scales. The key to this is a scale-breaking Morse theoretic argument (cf. Lemma 3.1), which allows us to show that the intermediate regions are topologically simple (i.e., planar domains). The key geometric input for this argument is the fact that the curvature is sufficiently small in a scale invariant sense in the intermediate region, a fact related to the half-space theorem for complete properly embedded minimal surfaces in R³.

This scale-breaking analysis of the intermediate regions forms the technical heart of our work. In ambient manifolds (M^n, g) with $4 \le n \le 7$, a similar argument works given appropriate modifications. Moreover, for n = 3, we show that uniform area bounds are unceessary for understanding the local picture of degeneration. Finally, everything we have discussed works for one-sided surfaces as well.

1.1. Applications of the local picture of degeneration. Thanks to our understanding of how embedded hypersurfaces with uniformly bounded index (and volume) can degenerate, we can prove various results along the lines of the general principle that (when the ambient dimension satisfies $3 \le n \le 7$) "embedded minimal hypersurfaces with uniformly bounded index behave qualitatively like embedded stable minimal hypersurfaces." We now discuss several results along these lines.

1.1.1. Finitely many diffeomorphism types. An easy application of curvature estimates for stable minimal hypersurfaces shows that for a closed Riemannian manifold (M^n, g) , $3 \le n \le 7$, there can be at most $N = N(M, g, \Lambda)$ distinct diffeomorphism types in the set of stable embedded minimal hypersurfaces with $\operatorname{vol}_g(\Sigma) \le \Lambda$. To see this, suppose that Σ_j is an infinite sequence of pairwise non-diffeomorphic embedded stable minimal hypersurfaces. Using the curvature estimates¹ established in [Sch83a, Ros06, SSY75, SS81] we see that if Σ_j is a sequence of embedded² stable minimal hypersurfaces, then there is C > 0 so that $|II_{\Sigma_j}|(x) \le C$ for all $x \in \Sigma_j$. Because $\operatorname{vol}_g(\Sigma_j) \le \Lambda$, by passing to

¹Note that the works [SSY75, SS81] only consider (embedded) two-sided stable hypersurfaces! However, using the fact that a properly embedded hypersurface in Euclidean space is two-sided, we can extend the curvature estimates to the one-sided case as well; see the proof of Lemma 2.4.

²Strictly speaking, embeddedness is not needed for this result in dimension n = 3. Note that it will be essential elsewhere in our work even when n = 3.

a subsequence we may find Σ_{∞} so that Σ_j converges locally smoothly to Σ_{∞} with finite multiplicity. Thus, for j sufficiently large we may construct a smooth covering map $\Sigma_j \to \Sigma_{\infty}$ with a uniformly bounded number of sheets. Because the Σ_j are assumed to be non-diffeomorphic, this easily yields a contradiction.

To try to extend this proof to the case of uniformly bounded index, we must contend with the possibility that the hypersurfaces have diverging curvature. Using our local picture of degeneration, we can deal with this possibility and show the following finiteness result.

Theorem 1.1. Fix (M^n, g) a closed Riemannian manifold, where $3 \le n \le 7$. Then there can be at most $N = N(M, g, \Lambda, I)$ distinct diffeomorphism types in the set of embedded minimal hypersurfaces $\Sigma \subset (M, g)$ with $index(\Sigma) \le I$ and $vol_g(\Sigma) \le \Lambda$.

In particular, for a closed three-manifold (M^3, g) , there is $r_0 = r_0(M, g, \Lambda, I)$ so that any embedded minimal surface Σ in (M^3, g) with $index(\Sigma) \leq I$ and $area_g(\Sigma) \leq \Lambda$ has $genus(\Sigma) \leq r_0$.

In a related direction, we can partially extend Ros's bounds [Ros06, Theorem 17] (see also [CM14]) to higher dimensions as follows.

Theorem 1.2. For $4 \le n \le 7$, there is $N = N(n, I, \Lambda) \in \mathbb{N}$ so that there are at most N mutually non-diffeomorphic complete embedded minimal hypersurfaces $\Sigma^{n-1} \subset \mathbb{R}^n$ with $\operatorname{index}(\Sigma) \le I$ and $\operatorname{vol}(\Sigma \cap B_R(0)) \le \Lambda R^{n-1}$ for all R > 0.

It would be interesting to understand how N depends on I and Λ .

1.1.2. Three-dimensional results. A well known compactness result for minimal surfaces in a fixed three-manifold is due to Choi–Schoen [CS85] who showed that for any sequence of minimal surfaces with bounded genus and area there is a subsequence converging to a smooth minimal surface (possibly with multiplicity). The convergence is moreover smooth away from finitely many points where curvature is concentrating.

This result has several important manifestations; for example, in a hyperbolic manifold, a genus bound for a minimal surface already implies an area bound by the Gauss equation and the Gauss-Bonnet formula. Moreover, in a three-manifold with positive Ricci curvature, the area of an embedded minimal surface can be bounded above in terms of its genus, by work of Choi–Wang [CW83] and Yang–Yau [YY80]. This bound and the non-existence of two-sided stable minimal hypersurfaces with ambient positive Ricci, shows that the Choi–Schoen compactness implies that the set of closed, embedded minimal surfaces with fixed genus in a three-manifold of positive Ricci curvature is compact in the smooth topology. In a general Riemannian three-manifolds, on the other hand, it is no longer possible to bound the area nor the index of an embedded minimal surface by the genus, even if one assumes positive scalar curvature, as it can be seen in examples constructed by Colding–De Lellis [CDL05].

However, we are able to show that in three-manifolds with positive scalar curvature, uniform index bounds do imply uniform area and genus bounds. This indicates that index bounds are not only very natural from the variational point of view, but they actually yield more control on the minimal surface than genus bounds. That such a result should hold follows again from our general principle that because this holds for embedded stable minimal surfaces, it should hold for an embedded minimal surface with bounded index. The corresponding result in the case of stable surfaces is a consequence of the fact that by work of Fischer-Colbrie–Schoen [FCS80] and Schoen–Yau [SY83], two-sided stable minimal surfaces in ambient manifolds with positive scalar curvature are S^2 , along with a geometric compactness argument based on the fact that S^2 is simply connected.

Theorem 1.3. Suppose that (M^3, g) is a closed three-manifold with positive scalar curvature. For $I \in \mathbb{N}$, there is $A_0 = A_0(M, g, I) < \infty$ and $r_0 = r_0(M, g, I)$ so that if $\Sigma \subset (M, g)$ is a connected, closed, embedded minimal surface with index $(\Sigma) \leq I$, then $\operatorname{area}_g(\Sigma) \leq A_0$ and $\operatorname{genus}(\Sigma) \leq r_0$.

Remark 1.4. Unfortunately, without any extra assumption, even the space of embedded stable minimal surfaces fails to be compact in general three-manifolds due to the failure of uniform area bounds. We discuss several examples in §1.2 below. In the converse direction, Ejiri–Micallef have shown [EM08] that for immersed minimal surfaces in a general three-manifold, uniform bounds on their area and genus imply uniform bounds on their index.

In a more technical direction, we remark that as a byproduct of the proof of Theorem 1.17, we obtain:

Theorem 1.5. Suppose that Σ_j is a sequence of embedded minimal surfaces in a threemanifold (M, g) with uniformly bounded index, i.e., $index(\Sigma_j) \leq I$ for some $I \in \mathbb{N}$. Then, after passing to a subsequence, Σ_j converges to a lamination \mathcal{L} away from at most I singular points. The lamination can be extended across these points.

Remark 1.6. We note that the fact that the limit lamination \mathcal{L} has removable singularities can be seen as a consequence of deep work by Meeks–Perez–Ros [MPR13], combined with our curvature estimates for Σ_j , which after passing to the limit, imply³ that $|\text{II}_{\mathcal{L}}|(x)d_g(x, \mathcal{B}_{\infty}) \leq C$ for $x \in \mathcal{L}$. Our proof of Theorem 1.5 however does not rely on the removable singularity results in [MPR13], and thus provides a self-contained proof that the limit lamination of a sequence of embedded closed minimal surfaces with bounded index has removable singularities.

Remark 1.7. We also remark that Theorem 1.17 and Corollary 1.19 provide an alternative approach to a recent result by Colding–Gabai, [CG14, Theorem 2.2] (cf. Remark 1.18). We will not reproduce the full statement here, but only note that it loosely says that a degenerating sequence of index-one embedded minimal surfaces will look like a small catenoid connected by large annular regions to the rest of the surface.

We remark that as consequence of the Theorems 1.1 and 1.3, we may easily deduce several compactness results. By a theorem of Colding–Minicozzi [CM00], the set of closed embedded minimal surfaces with uniformly bounded area and genus is finite in (M^3, g) , as long as g is "bumpy" in the sense of White [Whi15], i.e. g has the property

³Alternatively, [LZ15, (3.1)] establishes similar curvature estimates for the limit lamination; the exact form of estimates we establish here (before passing to the limit) are crucial for our proof of Theorem 1.17 in several other places.

that there are no immersed minimal submanifolds with non-zero Jacobi fields.⁴ Such metrics are "generic" by the main result in [Whi15]. Thus, we have:

Corollary 1.8. Suppose that (M^3, g) is a closed three-manifold with a bumpy metric of positive scalar curvature. For $I \in \mathbb{N}$, there are only finitely many closed, connected, embedded minimal surfaces Σ with index $(\Sigma) \leq I$.

Remark 1.9. A slightly different version of this corollary has recently been independently obtained by Carlotto [Carl5], assuming the ambient positive scalar curvature metric is bumpy in the sense of [Whi91] (i.e., there is no embedded (as opposed to immersed) minimal submanifold with a non-zero Jacobi field) but with the additional assumption that (M^3, q) contains no embedded minimal $\mathbb{R}P^2$.

Combining Theorem 1.3 with the work of Choi–Schoen [CS85], we also have:

Corollary 1.10. Suppose that (M^3, g) is a closed three-manifold with positive Ricci curvature. Then, for $I \in \mathbb{N}$, the set of closed, connected, embedded minimal surfaces Σ with index $(\Sigma) \leq I$ is compact in the smooth topology.

In particular, combined with the recent work of Marques–Neves [MN13] we obtain

Corollary 1.11. Suppose that (M^3, g) is a closed three-manifold with a bumpy metric of strictly positive Ricci curvature. Then, there exists a sequence of closed, embedded minimal surfaces Σ_j with index $(\Sigma_j) \to \infty$.

Remark 1.12. These last two corollaries have been recently proven by Li–Zhou [LZ15] by somewhat different arguments.

1.2. Counterexamples. Several examples show that the set of closed, embedded, stable minimal surfaces can fail to be compact, even if the metric is bumpy and even if we restrict only to the set of such surfaces with a fixed genus. The examples below show that the hypothesis in the applications discussed above cannot be significantly weakened.

Example 1.13. The simplest example of non-compactness occurs in the square threetorus $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$, equipped with the flat metric, as seen by choosing positive rational numbers $\theta_k \in \mathbb{Q}$ converging to an irrational number $\theta_\infty \in \mathbb{R} \setminus \mathbb{Q}$. Letting γ_k denote the simple closed geodesic in the two-torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ with slope θ_k , it is easy to see that $\Sigma_k := \gamma_k \times \mathbb{S}^1$ is an embedded stable minimal surface with $\operatorname{area}_g(\Sigma_k) \to \infty$. Note that the surfaces Σ_k limit to the lamination of \mathbb{T}^3 by a single plane $\gamma_\infty \times \mathbb{S}^1$ (where γ_∞ is the non-closed geodesic with slope θ_∞). Of course, the flat metric on \mathbb{T}^3 is manifestly non-bumpy, but it is relatively easy to see that for an arbitrary metric on \mathbb{T}^3 , we can minimize (by [SY79]) the *g*-area of immersions homotopic to the embedding of Σ_k into \mathbb{T}^3 and then argue (using [BW69, FHS83] and fundamental group considerations) that this yields a sequence of embedded, stable, minimal tori in (\mathbb{T}^3, g) with unbounded area.

⁴It seems to us that in general, the notion of "bumpy" from [Whi15], rather than the notion from [Whi91] is necessary to deal with the possibility of a one-sided limit in the proof of [CM00]. See also [Car15, Remark 3.1].

One might hope that the torus \mathbb{T}^3 is somehow special in the previous example. However, the following results show that for *any* closed three-manifold, it is not possible to use bumpiness (or more generally, any "generic" property which is satisfied by a C^2 dense set of metrics) to prove area bounds for embedded stable minimal surfaces (even assuming fixed genus).

Example 1.14. Fix a Riemannian three-manifold M. Work of Colding–Minicozzi [CM00] shows that there is a C^2 -open set of metrics g on M so that there is a sequence of embedded stable minimal tori Σ_j with $\operatorname{area}_g(\Sigma_j) \to \infty$. Indeed, for any domain in M of the form $\Omega = S \times \mathbb{S}^1$ where S is a disk with three holes removed (i.e., Ω is a solid torus with three holes removed, which obviously exists in any coordinate chart), they show that if Ω has strictly mean convex boundary with respect to a metric g, then there exists such a sequence of tori in $\Omega \subset (M, g)$. Their construction relies on an idea of "looping" tori around the holes; see [Kra09, Figure 3.2.3] for a nice illustration. These examples were subsequently extended by Dean [Dea03] and Kramer [Kra09] to give examples of sequences of embedded, stable, minimal surfaces with unbounded area, with any fixed genus.

Example 1.15. An even more extreme example similar to Example 1.14 but with a more complicated looping scheme was given by Colding–Hingston [CH06], who in a C^2 -open set of metrics on any three-manifold, construct a sequence of stable tori with unbounded area whose limit lamination has surprising behavior, in particular non-closed limit leaves.

In a more topological vein, we have the following examples of embedded minimal surfaces with bounded index (in fact stable) but unbounded genus (and hence area).

Example 1.16. For Σ_r the closed oriented surface of genus r > 1, Jaco proved [Jac70] that $M_r := \Sigma_r \times \mathbb{S}^1$ admits a sequence of incompressible surfaces with unbounded genus. For any Riemannian metric g on M_r , we may minimize area using [SY79] and see that the resulting stable minimal surface is embedded (after passing to a one-sided quotient, if necessary) by [FHS83]. It is clear that these minimal surfaces must have unbounded genus.

1.3. Precise statement of degeneration and surgery results in 3-dimensions. We now state our main results in three-dimensions.

Theorem 1.17 (Local picture of degeneration). There are functions m(I) and r(I) with the following property. Fix a closed three-manifold (M^3, g) and a natural number $I \in \mathbb{N}$. Then, if $\Sigma_i \subset (M, g)$ is a sequence of closed embedded minimal surfaces with

 $\operatorname{index}(\Sigma_j) \leq I,$

then after passing to a subsequence, there is C > 0 and a finite set of points $\mathcal{B}_j \subset \Sigma_j$ with cardinality $|\mathcal{B}_j| \leq I$ so that the curvature of Σ_j is uniformly bounded away from the set \mathcal{B}_j , i.e.,

 $|\mathrm{II}_{\Sigma_j}|(x)\min\{1, d_g(x, \mathcal{B}_j)\} \le C,$

but not at \mathcal{B}_j , i.e.,

 $\liminf_{j\to\infty}\min_{p\in\mathcal{B}_j}|\mathrm{II}_{\Sigma_j}|(p)=\infty.$

Passing to a further subsequence, the points \mathcal{B}_j converge to a set of points \mathcal{B}_{∞} and the surfaces Σ_j converge locally smoothly, away from \mathcal{B}_{∞} , to some lamination $\mathcal{L} \subset M \setminus \mathcal{B}_{\infty}$. The lamination has removable singularities, i.e., there is a smooth lamination $\widetilde{\mathcal{L}} \subset M$ so that $\mathcal{L} = \widetilde{\mathcal{L}} \setminus \mathcal{B}_{\infty}$. Moreover, there exists $\varepsilon_0 > 0$ smaller than the injectivity radius of (M, g) so that \mathcal{B}_{∞} is $4\varepsilon_0$ -separated and for any $\varepsilon \in (0, \varepsilon_0]$, taking j sufficiently large guarantees that

- (1) Writing Σ'_j for the components of $\Sigma_j \cap B_{2\varepsilon}(\mathcal{B}_{\infty})$ containing at least one point from \mathcal{B}_j , no component of Σ'_j is a topological disk, so we call Σ'_j the "neck components." They have the following additional properties:
 - (1.a) The surface Σ'_j intersects $\partial B_{\varepsilon}(\mathcal{B}_{\infty})$ transversely in at most m(I) simple closed curves.
 - (1.b) Each component of Σ'_i is unstable.
 - (1.c) The genus⁵ of Σ'_i is bounded above by r(I).
 - (1.d) The area of Σ'_i is uniformly bounded, i.e.,

$$\limsup_{j \to \infty} \operatorname{area}_g(\Sigma'_j) \le 2\pi m(I)\varepsilon^2(1+o(\varepsilon))$$

(2) Writing Σ_j'' for the components of $\Sigma_j \cap B_{2\varepsilon}(\mathcal{B}_{\infty})$ that do not contain any points in \mathcal{B}_j , each component of Σ_j'' is a topological disk, so we call Σ_j'' the "disk components." Moreover, we have the following additional properties

(2.a) The curvature of $\Sigma_{i}^{\prime\prime}$ is uniformly bounded, i.e.,

$$\limsup_{j \to \infty} \sup_{x \in \Sigma_j''} |\mathrm{II}_{\Sigma_j}|(x) < \infty$$

(2.b) Each component of Σ''_j has area uniformly bounded above by $2\pi\varepsilon^2(1 + o(\varepsilon))$.

As is clear from the proof, it would be possible to give explicit bounds for m(I) and r(I), if one desired.

Remark 1.18. As remarked above, a similar description in the special case of index one surfaces was recently obtained by Colding–Gabai [CG14, Theorem 2.2]. However, our proof differs from theirs (even in the index one case) in how we transfer topological information between scales. Additionally, the higher index case introduces serious technical difficulties, due to the possibility of simultaneous concentration at multiple scales.

A key application of Theorem 1.17 is a prescription for performing "surgery" on a sequence of bounded index minimal surfaces so that their curvature remains bounded, while only changing the topology and geometry in a controllable way.

Corollary 1.19 (Controlled surgery). There exist functions $\tilde{r}(I)$ and $\tilde{m}(I)$ with the following property. Fix a closed three-manifold (M^3, g) and suppose that $\Sigma_j \subset (M^3, g)$

⁵See Definition A.2.

is a sequence of closed embedded minimal surfaces with

$$\operatorname{index}(\Sigma_i) \leq I.$$

Then, after passing to a subsequence, there is a finite set of points $\mathcal{B}_{\infty} \subset M$ with $|\mathcal{B}_{\infty}| \leq I$ and $\varepsilon_0 > 0$ smaller than the injectivity radius of (M,g) so \mathcal{B}_{∞} is $4\varepsilon_0$ -separated, and so that for $\varepsilon \in (0, \varepsilon_0]$, if we take j sufficiently large then there exists embedded surfaces $\tilde{\Sigma}_j \subset (M^3, g)$ satisfying:

- (1) The new surfaces $\widetilde{\Sigma}_i$ agree with Σ_i outside of $B_{\varepsilon}(\mathcal{B}_{\infty})$.
- (2) The components of $\Sigma_j \cap B_{\varepsilon}(\mathcal{B}_{\infty})$ that do not intersect the spheres $\partial B_{\varepsilon}(\mathcal{B}_{\infty})$ transversely and those that are topological disks appear in $\widetilde{\Sigma}_j$ without any change.
- (3) The curvature of $\widetilde{\Sigma}_i$ is uniformly bounded, i.e.

$$\limsup_{j \to \infty} \sup_{x \in \widetilde{\Sigma}_j} |\mathrm{II}_{\widetilde{\Sigma}_j}|(x) < \infty.$$

- (4) Each component of $\widetilde{\Sigma}_j \cap B_{\varepsilon}(\mathcal{B}_{\infty})$ which is not a component of $\Sigma_j \cap B_{\varepsilon}(\mathcal{B}_{\infty})$ is a topological disk of area at most $2\pi\varepsilon^2(1+o(\varepsilon))$.
- (5) The genus drops in controlled manner, i.e.,

$$\operatorname{genus}(\Sigma_j) - \tilde{r}(I) \leq \operatorname{genus}(\widetilde{\Sigma}_j) \leq \operatorname{genus}(\Sigma_j).$$

(6) The number of connected components increases in a controlled manner, i.e.,

$$|\pi_0(\Sigma_j)| \le |\pi_0(\widetilde{\Sigma}_j)| \le |\pi_0(\Sigma_j)| + \widetilde{m}(I).$$

(7) While $\widetilde{\Sigma}_j$ is not necessarily minimal, it is asymptotically minimal in the sense that $\lim_{j\to\infty} \|H_{\widetilde{\Sigma}_j}\|_{L^{\infty}(\widetilde{\Sigma}_j)} = 0.$

The new surfaces $\widetilde{\Sigma}_j$ converge locally smoothly to the smooth minimal lamination $\widetilde{\mathcal{L}}$ from Theorem 1.17.

Remark 1.20. The strategy we use to prove Theorem 1.17 can be extended to a higher dimensional setting (assuming a uniform volume bound). Certain aspects of the local structure change; in particular, due to the failure of the half-space theorem in higher dimensions, the separation of sheets into "neck regions" and "disk regions" does not occur in the same way as in three dimensions. As such, we will not attempt to formulate a higher dimensional version of Theorem 1.17 or Corollary 1.19, but from the proof of Theorems 1.1 and 1.2 it is clear that the general picture described in the introduction holds.

1.4. Related Results. As remarked above, Li–Zhou [LZ15] have proven compactness results for embedded minimal surfaces with bounded index in a three-manifold with a metric of positive Ricci curvature. This was preceded by the higher dimensional (i.e., allowing the ambient manifold to be *n*-dimensional for $3 \le n \le 7$) result of Sharp [Sha15], showing that for a metric of positive Ricci curvature, the space of embedded minimal surfaces with uniformly bounded area and index is compact.

After this paper was completed, we were informed by Carlotto that he had independently arrived at a proof of a slightly different version of Corollary 1.8. His paper [Car15] appeared at essentially the same time as ours.

These works mainly focus on properties of limits of surfaces with uniformly bounded index, rather than the way in which such surfaces degenerate. As such, their arguments are of a rather different nature than those in this paper.

The version of this article originally posted to the arXiv only discussed the case of ambient three manifolds. After we had completed extending our work to higher dimensions, but before we had updated the arXiv version, we received a preprint [BS15] from Buzano and Sharp containing an alternative approach to the topological bounds in higher dimensions. Their work recovers Theorem 1.1 via a rather different approach.

Finally, we refer to the works of Ros [Ros95] and Traizet [Tra04] studying of how complete embedded minimal surfaces in \mathbb{R}^3 with bounded total curvature degenerate. Some parallels can be drawn between their results and Theorem 1.17, but our work takes a different technical approach due to the precise behavior of the index.

1.5. Outline of the paper. In Section 2, we make several preliminary definitions and prove curvature bounds away from finitely many points for hypersurfaces with bounded index. Section 3 contains a key topological result allowing us to control the topology of the "intermediate regions", assuming a curvature bound of the appropriate form. We establish the local picture of degeneration for in three-manifolds in Section 4 and the surgery result in Section 5. These results then allow us to establish the three-dimensional compactness results in Section 6. Section 7 contains the proof of the higher dimensional results. Appendix A contains a discussion of the non-orientable genus as well as the genus of surfaces with boundary. Appendix B recalls certain facts about finite index surfaces in \mathbb{R}^3 . In Appendix C we provide proofs of several removable singularity results. Finally, Appendix D contains examples to illustrate that the various forms of degeneration discussed in the proof of Propositions 4.2 and 4.3 can in fact occur.

1.6. Acknowledgements. We are grateful to Dave Gabai for several illuminating conversations. We thank Nick Edelen for several useful comments and Alessandro Carlotto for pointing out a mistaken reference in an earlier version of Corollary 1.8. O.C. would like to acknowledge Jacob Bernstein for a helpful discussion regarding removable singularity results and Sander Kupers for kindly answering some questions about topology. He was partially supported by an NSF Graduate Research Fellowship DGE-1147470 during the time which most of this work was undertaken, and also acknowledges support by the EPSRC Programme Grant, Singularities of Geometric Partial Differential Equations number EP/K00865X/1. D.K. was partially supported by an NSF Post-doctoral Fellowship DMS-1401996. D.M. would like to acknowledge Rick Schoen for several helpful conversations. He was partially supported by NSF grant DMS-1512574 during the completion of this work.

2. Preliminaries

2.1. Definitions and basic notation. Let Σ be a closed embedded minimal hypersurface in (M, g). Recall, whether Σ is one-sided or two-sided, the *Morse index* of Σ , henceforth denoted by $index(\Sigma)$, is defined as the number of negative eigenvalues of the quadratic form associated to second variation of area:

$$Q(v,v) := \int_{\Sigma} |\nabla^{\perp} v|^2 - |\mathrm{II}_{\Sigma}|^2 |v|^2 - \mathrm{Ric}_M(v,v) \, d\Sigma,$$

where v is a section of the normal bundle of Σ in M, ∇^{\perp} the induced connection, and Π_{Σ} the second fundamental form of Σ . Whenever Σ is two-sided, the Morse index is equal to the number of negative eigenvalues of the associated Jacobi operator $\Delta_{\Sigma} + |\Pi_{\Sigma}|^2 + \operatorname{Ric}(\nu, \nu)$ acting on smooth functions $\varphi \in C^{\infty}(\Sigma)$. If Σ is one-sided, however, we consider the orientable double cover $\hat{\Sigma} \to \Sigma$ and the corresponding change of sheets involution of $\tau : \hat{\Sigma} \to \hat{\Sigma}$, which must satisfy $\nu \circ \tau = -\nu$ for any choice on unit normal vector ν for $\hat{\Sigma}$, then the Morse index of Σ is then equal to the negative eigenvalues of the operator $\Delta_{\hat{\Sigma}} + |\Pi_{\hat{\Sigma}}|^2 + \operatorname{Ric}(\nu, \nu)$ over the space of smooth functions $\varphi \in C^{\infty}(\hat{\Sigma})$ satisfying $\varphi \circ \tau = -\varphi$.

We will be dealing with sequences of embedded minimal surfaces without area bounds in three manifolds and so it will be convenient to consider minimal laminations: A closed set \mathcal{L} in M^3 is called a *minimal lamination* if \mathcal{L} is the union of pairwise disjoint, connected, injectively immersed minimal surfaces, called *leaves*. For each point $x \in M^3$, we require the existence of a neighborhood $x \in \Omega$ and a $C^{0,\alpha}$ local coordinate chart $\Phi : \Omega \to \mathbb{R}^3$ under which image the leaves of \mathcal{L} pass through in slices of the form $\mathbb{R}^2 \times \{t\} \cap \Phi(\Omega)$.

All distance functions considered in our work will be induced by some ambient metric, and we denote by d_h the distance function induced by the metric h. Given a closed set S, we let $d_h(\cdot, S)$ denote the distance to S with respect to the metric h. If S is a finite set of points, |S| will denote its cardinality, and for some $\delta > 0$, we say that S is δ -separated if $d_h(x, S \setminus \{x\}) > \delta$ for every $x \in S$. We will also consider metric balls and write, as usual, $B_r(p)$ to denote the ball of radius r > 0 centered at p. If a finite set of points is δ -separated and $\delta > r > 0$, then the set of points within distance at most rfrom S forms a union of disjoint balls, which we will be denote by $B_r(S)$ (Note: we are omit the dependence of the ambient metric in our notation for B_r as it should always be clear in the context in which is being used).

2.1.1. Smooth blow-up sets. The following definition turns out to be quite convenient in the sections to come. Suppose that $(M_j, g_j, 0_j)$ is a sequence⁶ of complete pointed Riemannian manifolds which are converging in the pointed Cheeger–Gromov sense to $(M_{\infty}, g_{\infty}, 0_{\infty})$. Suppose that Σ_j is a sequence of embedded minimal hypersurfaces in (M_j, g_j) . A sequence of finite sets of points $\mathcal{B}_j \subset \Sigma_j$ is said to be a sequence of smooth blow-up sets if:

(1) The set \mathcal{B}_i remains a finite distance from the basepoint 0_i , i.e.

$$\limsup_{j \to \infty} \max_{p \in \mathcal{B}_j} d_{g_j}(p, 0_j) < \infty.$$

⁶In practice, either $(M_j, g_j, 0_j)$ will be a fixed (independently of j) compact manifold or $(M_{\infty}, g_{\infty}, 0_{\infty})$ will be Euclidean space.

(2) If we set $\lambda_j(p) := |II_{\Sigma_j}|(p)$ for $p \in \mathcal{B}_j$, then the curvature of Σ_j blows up at each point in \mathcal{B}_j , i.e.,

$$\liminf_{j \to \infty} \min_{p \in \mathcal{B}_j} \lambda_j(p) = \infty.$$

(3) If we choose a sequence of points $p_j \in \mathcal{B}_j$, then after passing to a subsequence, the rescaled surfaces $\overline{\Sigma}_j := \lambda_j(p_j)(\Sigma_j - p_j)$ converge locally smoothly to a complete, non-flat, embedded minimal surface $\overline{\Sigma}_{\infty} \subset \mathbb{R}^n$ without boundary, satisfying

$$|\mathrm{II}_{\overline{\Sigma}_{\infty}}|(x) \le |\mathrm{II}_{\overline{\Sigma}_{\infty}}|(0),$$

for all $x \in \mathbb{R}^n$.

(4) The blow-up points do not appear in the blow-up limit of the other points, i.e.,

$$\liminf_{\substack{j \to \infty}} \min_{\substack{p, q \in \mathcal{B}_j \\ p \neq q}} \lambda_j(p) d_{g_j}(p, q) = \infty.$$

2.2. Curvature estimates and index concentration. Recall that Schoen [Sch83a] has proven that two-sided stable minimal surfaces in a three-manifold have uniformly bounded curvature. More recently, the two-sided hypothesis was shown to be unnecessary by Ros [Ros06]. In particular, we have:

Theorem 2.1 ([Sch83a, Ros06]). Fix (M^3, g) a closed three-manifold. There is C = C(M, g) so that if $\Sigma \subset (M, g)$ is a compact stable minimal surface, then

$$|\mathrm{II}_{\Sigma}|(x)\min\{1, d_g(x, \partial \Sigma)\} \leq C$$

for all $x \in \Sigma$.

Here, we show that sequence of embedded minimal surfaces of bounded index have curvature bounds away from at most finitely many points. This can be thought of as a generalization of Schoen and Ros's curvature estimates for stable minimal surfaces to the case of finite Morse index. The proof by induction is most convenient if we prove a more general bound for surfaces with boundary.

Lemma 2.2. Fix (M^3, g) a closed three-manifold and $I \in \mathbb{N}$. Suppose that $\Sigma_j \subset (M, g)$ is a sequence of compact embedded minimal surfaces with $index(\Sigma_j) \leq I$. Then, after passing to a subsequence, there exist C > 0 and a sequence of smooth blow-up sets $\mathcal{B}_j \subset \Sigma_j$ with $|\mathcal{B}_j| \leq I$, so that

$$|\mathrm{II}_{\Sigma_i}|(x)\min\{1, d_q(x, \mathcal{B}_j \cup \partial \Sigma_j)\} \le C.$$

for all $x \in \Sigma_j$.

Proof. We prove this by induction on I. When I = 0, the surface Σ is stable, so the statement is exactly the curvature estimates discussed in Theorem 2.1.

For I > 0, consider $\Sigma_j \subset (M, g)$ with $index(\Sigma_j) \leq I$. By passing to a subsequence, we may assume that

$$\rho_j := \sup_{x \in \Sigma_j} |\mathrm{II}_{\Sigma_j}|(x) \min\{1, d_g(x, \partial \Sigma_j)\} \to \infty.$$

If we cannot find such a subsequence, it is easy to see that the curvature estimates hold with $\mathcal{B}_{i} = \emptyset$.

A standard point picking argument allows us to find $\tilde{p}_j \in \Sigma_j$ so that for $\lambda_j = |II_{\Sigma_j}|(\tilde{p}_j) \to \infty$, the rescaled surfaces

$$\overline{\Sigma}_j := \lambda_j (\Sigma_j - \widetilde{p}_j)$$

converge locally smoothly, after passing to a subsequence, to an embedded⁷ minimal surface in \mathbb{R}^3 , $\widehat{\Sigma}_{\infty}$, of index at most *I* and with no boundary, so that

$$|\mathrm{II}_{\widehat{\Sigma}_{\infty}}|(x) \le |\mathrm{II}_{\widehat{\Sigma}_{\infty}}|(0) = 1.$$

For the reader's convenience, we recall the point picking argument at the end of the proof.

Because $\widehat{\Sigma}_{\infty}$ is non-flat, there is some radius $\widehat{R} > 0$ so that $\widehat{\Sigma}_{\infty} \cap B_{\widehat{R}}(0)$ has nonzero index, $\widehat{\Sigma}_{\infty} \setminus B_{\widehat{R}}(0)$ is stable, and $\widehat{\Sigma}_{\infty}$ intersects $\partial B_R(0)$ transversely. Moreover, taking \widehat{R} larger if necessary, we may arrange that all of these properties are satisfied in addition to

$$(2.1) |II_{\widehat{\Sigma}_{\infty}}|(x) \le \frac{1}{4}.$$

for $x \in \widehat{\Sigma}_{\infty} \setminus B_{\widehat{R}}(0)$.

We define $\widetilde{\Sigma}_j := \Sigma_j \setminus B_{\widehat{R}/\lambda_j}(\widetilde{p}_j)$. For j large, this ball cannot intersect the boundary of Σ_j (by the choice of \widetilde{p}_j and because $\rho_j \to \infty$) and $\partial B_{\widehat{R}/\lambda_j}(\widetilde{p}_j)$ intersects Σ_j transversely. Thus, $\widetilde{\Sigma}_j$ is a smooth compact minimal surface with smooth, compact boundary

$$\partial \widetilde{\Sigma}_j = \partial \Sigma_j \cup (\partial B_{\widehat{R}/\lambda_j}(\widetilde{p}_j) \cap \Sigma_j).$$

For j large, $\operatorname{index}(\widetilde{\Sigma}_j) \leq I - 1$. By the inductive hypothesis, passing to a subsequence, there is a sequence of smooth blow-up sets $\widetilde{\mathcal{B}}_j \subset \Sigma_j$ with $|\widetilde{\mathcal{B}}_j| \leq I - 1$ and a constant \widetilde{C} (independent of j) so that

(2.2)
$$|\mathrm{II}_{\widetilde{\Sigma}_j}|(x)\min\{1, d_g(x, \widetilde{\mathcal{B}}_j \cup \partial \widetilde{\Sigma}_j)\} \le \widetilde{C}.$$

We claim that $\mathcal{B}_j := \widetilde{\mathcal{B}}_j \cup {\widetilde{p}_j}$ is a sequence of smooth blow-up sets. The only thing we must check is that none of the points in $\widetilde{\mathcal{B}}_j$ can appear in the blow-up around \widetilde{p}_j and vice versa (in particular, this guarantees that rescaling Σ_j around points in $\widetilde{\mathcal{B}}_j$ still yields a smooth limit). First, suppose that

$$\liminf_{j\to\infty}\min_{\widetilde{r}\in\widetilde{\mathcal{B}}_j}\lambda_j d_{g_j}(\widetilde{r},\widetilde{p}_j)<\infty,$$

where we recall that $\lambda_j = |\mathrm{II}_{\Sigma_j}|(\widetilde{p}_j)$. Assume that for j fixed, the minimum is attained at $\widetilde{r}_j \in \widetilde{\mathcal{B}}_j$. By choice of \widehat{R} (specifically (2.1)) we see that after passing to a subsequence,

$$\eta_j := |\mathrm{II}_{\Sigma_j}|(\widetilde{r}_j) \le \frac{1}{2}|\mathrm{II}_{\Sigma_j}|(\widetilde{p}_j) = \frac{1}{2}\lambda_j.$$

Thus, we have reduced to the other possibility, i.e.

$$\liminf_{j \to \infty} \eta_j d_{g_j}(\widetilde{r}_j, \widetilde{p}_j) < \infty.$$

⁷Usually, the blow-up limit $\hat{\Sigma}_{\infty}$ would only be injectively immersed. Here, because it has finite index and no boundary, it is embedded by Theorem B.1.

However, this is a contradiction, as the blow-up of $\widetilde{\Sigma}_j$ around \widetilde{r}_j has no boundary.

Now, suppose that there is $z_j \in \Sigma_j$ so that

$$\limsup_{j \to \infty} |\mathrm{II}_{\Sigma_j}|(z_j) \min\{1, d_{g_j}(z_j, \mathcal{B}_j \cup \partial \Sigma_j)\} = \infty.$$

Combined with (2.2) and choice of \widetilde{p}_j , we may pass to a subsequence with $z_j \in \widetilde{\Sigma}_j$ and

$$d_{g_j}(z_j, \mathcal{B}_j \cup \partial \Sigma_j) = d_{g_j}(z_j, \widetilde{p}_j) \to 0,$$

$$d_{g_j}(z_j, \widetilde{\mathcal{B}}_j \cup \partial \widetilde{\Sigma}_j) = d_{g_j}(z_j, \widetilde{p}_j) - \frac{\widehat{R}}{\lambda_i}.$$

Because $\widehat{\Sigma}_{\infty}$ has bounded curvature, we see that z_j cannot appear in the blow-up around \widetilde{p}_j , i.e.,

$$\liminf_{j \to \infty} \lambda_j d_{g_j}(z_j, \widetilde{p}_j) = \infty.$$

Thus,

$$\limsup_{j \to \infty} |\mathrm{II}_{\Sigma_j}|(z_j) \frac{\widehat{R}}{\lambda_j} \le \limsup_{j \to \infty} \frac{\widetilde{C}\widehat{R}}{\lambda_j d_{g_j}(z_j, \widetilde{\mathcal{B}}_j \cup \partial \widetilde{\Sigma}_j)} = 0.$$

Combined with (2.2), this implies that

$$\widetilde{C} \geq \limsup_{j \to \infty} |\mathrm{II}_{\Sigma_j}|(z_j) \min\{1, d_{g_j}(z_j, \widetilde{\mathcal{B}}_j \cup \partial \widetilde{\Sigma}_j)\} = \infty,$$

a contradiction. This completes the proof.

Finally, we recall the point-picking argument used above to construct $\hat{\Sigma}_{\infty}$. Choose $\tilde{q}_j \in \Sigma_j$ so that

$$II_{\Sigma_j}|(\tilde{q}_j)\min\{1, d_g(\tilde{q}_j, \partial \Sigma_j)\} = \rho_j \to \infty$$

and set $r_j = |II_{\Sigma_j}|(\tilde{q}_j)^{-\frac{1}{2}}$. Then, choose $\tilde{p}_j \in \Sigma_j \cap B_{r_j}(\tilde{q}_j)$ so that

$$|\mathrm{II}_{\Sigma_j}|(\widetilde{p}_j)d_g(\widetilde{p}_j,\partial B_{r_j}(\widetilde{q}_j)) = \max_{x \in \Sigma_j \cap B_{r_j}(\widetilde{q}_j)} |\mathrm{II}_{\Sigma_j}|(x)d_g(x,\partial B_{r_j}(\widetilde{q}_j)).$$

Let $R_j = d_g(\widetilde{p}_j, \partial B_{r_j}(\widetilde{q}_j))$. Because $d_g(x, \partial B_{R_j}(\widetilde{p}_j)) \leq d_g(x, \partial B_{r_j}(\widetilde{q}_j))$ for $x \in B_{R_j}(\widetilde{p}_j)$, we find that

$$|\mathrm{II}_{\Sigma_j}|(\widetilde{p}_j)d_g(\widetilde{p}_j,\partial B_{R_j}(\widetilde{p}_j)) = \max_{x \in \Sigma_j \cap B_{R_j}(\widetilde{p}_j)} |\mathrm{II}_{\Sigma_j}|(x)d_g(x,\partial B_{R_j}(\widetilde{p}_j)).$$

Note that $|II_{\Sigma_j}|(\widetilde{p}_j)R_j \geq |II_{\Sigma_j}|(\widetilde{q}_j)r_j \to \infty$. The rescaled surfaces $\overline{\Sigma}_j$ satisfy

$$|\mathrm{II}_{\overline{\Sigma}_{j}}|(x)d_{\overline{g}_{j}}(x,\partial B_{\lambda_{j}R_{j}}(0)) \leq \lambda_{j}R_{j},$$

for $x \in \overline{\Sigma}_j \cap B_{\lambda_j R_j}(0)$. Thus, if $x \in \overline{\Sigma}_j$ lies in a given compact set of \mathbb{R}^3 , then

$$|\mathrm{II}_{\overline{\Sigma}_j}|(x) \le \frac{\lambda_j R_j}{\lambda_j R_j - d_{\overline{g}_j}(x,0)} \to 1 = |\mathrm{II}_{\overline{\Sigma}_j}|(0)$$

as $j \to \infty$. By construction, we see that $d_{\overline{g}_j}(0, \partial \overline{\Sigma}_j) \to \infty$. Passing to a subsequence, we may take a smooth limit of $\lambda_j(\Sigma_j - \widetilde{p}_j)$ to find a complete, non-flat, embedded minimal surface $\widehat{\Sigma}_{\infty}$ in \mathbb{R}^3 of index at most I and with no boundary, completing the point picking argument. **Corollary 2.3.** For (M^3, g) and $I \in \mathbb{N}$, if $\Sigma_j \subset (M, g)$ is a sequence of closed embedded minimal surfaces with $index(\Sigma_j) \leq I$, after passing to a subsequence, there is C > 0and a sequence smooth blow-up sets $\mathcal{B}_j \subset \Sigma_j$ with $|\mathcal{B}_j| \leq I$, so that

$$|\mathrm{II}_{\Sigma_j}|(x)d_g(x,\mathcal{B}_j) \le C,$$

for all $x \in \Sigma_j$.

In higher dimensions, we similarly have the following curvature estimates.

Lemma 2.4. Fix, for $4 \le n \le 7$, a closed n-dimensional manifold (M^n, g) , as well as $\Lambda > 0$ and $I \in \mathbb{N}$. Suppose that $\Sigma_j \subset (M, g)$ is a sequence of compact embedded minimal hypersurfaces with $\operatorname{vol}_g(\Sigma_j) \le \Lambda$ and $\operatorname{index}(\Sigma_j) \le I$. Then, after passing to a subsequence, there exists C > 0 and a sequence of smooth blow-up sets $\mathcal{B}_j \subset \Sigma_j$ with $|\mathcal{B}_j| \le I$, so that

$$|\mathrm{II}_{\Sigma_i}|(x)\min\{1, d_q(x, \mathcal{B}_i \cup \partial \Sigma_i)\} \le C.$$

for all $x \in \Sigma_j$.

Proof. The argument is similar to Lemma 2.2, so we will be brief. By [Sim68, SSY75, SS81], if $\hat{\Sigma}$ is an embedded, two-sided stable minimal hypersuface in \mathbb{R}^n (for $4 \leq n \leq$ 7), with Euclidean volume growth, i.e. $\lim_{R\to\infty} R^{1-n} \operatorname{vol}(\hat{\Sigma} \cap B_R) < \infty$, then it is a finite union of finitely many parallel planes. Because a complete properly embedded hypersurface in \mathbb{R}^n is two-sided (cf. [Sam69]) and an embedded hypersurface in \mathbb{R}^n with bounded second fundamental form and Euclidean volume growth is easily seen to be properly embedded, we obtain the following Bernstein-type result: if $\hat{\Sigma}$ is an embedded stable minimal hypersurface in \mathbb{R}^n (for $4 \leq n \leq 7$) with Euclidean volume growth and bounded second fundamental form, then it is the union of finitely many parallel planes. In particular, we do not need to assume a priori that $\hat{\Sigma}$ is two-sided.

From this, we obtain the claim when I = 0. Indeed, if it were false, we could find a sequence of compact embedded stable minimal hypersurfaces $\Sigma_j \subset (M^n, g)$ with $\operatorname{vol}_g(\Sigma_j) \leq \Lambda$. The point picking argument used above then produces a non-flat, embedded stable minimal hypersurface in \mathbb{R}^n with Euclidean volume growth (by the volume bounds and monotonicity formula) and uniformly bounded second fundamental form. This contradicts the above observation.

More generally, assume that the result fails for some fixed index bound I for a sequence Σ_j . Then, as in the proof of Lemma 2.2, we may find $\tilde{p}_j \in \Sigma_j$ so that for $\lambda_j = |\text{II}_{\Sigma_j}|(\tilde{p}_j)$, the rescaled surfaces

$$\overline{\Sigma}_j := \lambda_j (\Sigma_j - \widetilde{p}_j)$$

converge locally smoothly, after passing to a subsequence, to an embedded stable minimal hypersurface $\widehat{\Sigma}_{\infty}$ in \mathbb{R}^n with Euclidean area growth with

$$|\mathrm{II}_{\widehat{\Sigma}_{\infty}}|(x) \le |\mathrm{II}_{\widehat{\Sigma}_{\infty}}|(0) = 1.$$

Unlike the case when n = 3, it is possible that $\hat{\Sigma}_{\infty}$ has multiple components.⁸ However, the monotonicity formula guarantees that the number of components is bounded. Thanks to the above observation (implying that the index of $\hat{\Sigma}_{\infty}$ is non-zero) and the

⁸Recall that the half-space theorem fails for $n \ge 4$. For example, for $n \ge 4$, a catenoid in \mathbb{R}^n is bounded between two parallel planes.

fact that there are only finitely many components, we may choose \hat{R} exactly as in the proof of Lemma 2.4. The rest of the proof proceeds by removing $B_{\hat{R}/\lambda_j}(p_j)$ from Σ_j and using the inductive step, exactly as in Lemma 2.2.

3. ANNULAR DECOMPOSITION FROM CURVATURE ESTIMATES

The following lemma is a generalization of [Whi87, p. 251] (see also [MPR13, Lemma 4.1]). It will play a crucial role in later arguments, allowing us to transmit topological information between different scales.

Lemma 3.1 (Annular decomposition). There is $0 < \tau_0 < \frac{1}{2}$ with the following property. Assume that g is a Riemannian metric on $\{|x| \leq 2\} \subset \mathbb{R}^n$ which is sufficiently smoothly close to $g_{\mathbb{R}^n}$. Suppose that $\Sigma \subset B_1(0)$ is a properly embedded hypersurface with $\partial \Sigma \subset \partial B_1(0)$. Assume that for some $\tau \leq \tau_0$ and $p \in B_{\tau_0}(0)$, we have:

- (1) Each component of Σ intersects $B_{\tau}(p)$.
- (2) The hypersurface Σ intersects $\partial B_{\tau}(p)$ transversely in m manifolds diffeomorphic to \mathbb{S}^{n-2} with the standard smooth structure.
- (3) The curvature of Σ satisfies $|II_{\Sigma}|(x)d_g(x,p) \leq \frac{1}{4}$ for all $x \in \Sigma \setminus B_{\tau}(p)$.

Then, Σ intersects $\partial B_1(0)$ transversely in m manifolds diffeomorphic to the standard \mathbb{S}^{n-2} and each component of $\Sigma \cap \left(\overline{B_1(0)} \setminus B_\tau(p)\right)$ is diffeomorphic to $\mathbb{S}^{n-2} \times [0,1]$ with the standard smooth structure.

Proof. As long as g is sufficiently close to $g_{\mathbb{R}^n}$, working in normal coordinates around p, a computation as in [HI01, pp. 417–8] shows that the third hypothesis implies that the curvature of Σ with respect to $g_{\mathbb{R}^n}$ satisfies

$$|\mathrm{II}_{\Sigma}^{\mathbb{R}^n}|(x)d_{\mathbb{R}^n}(x,p) \leq \frac{1}{2}.$$

Hence, it is not hard to check that it is suffices to take $g = g_{\mathbb{R}^n}$.

Choose $\chi \in C_c^{\infty}([0,1))$ a smooth positive cutoff function so that $\chi(r) \in [0,1]$, $\chi(r) = 1$ for $r \leq \frac{1}{4}$ and $\chi(r) = 0$ for r sufficiently close to 1. We will take $\tau > 0$ sufficiently small based on this fixed cutoff function. Consider the function

$$f(x) = d_{\mathbb{R}^n}(x,p)^2 \chi(d_{\mathbb{R}^n}(0,x)^2) + d_{\mathbb{R}^n}(x,0)^2 (1 - \chi(d_{\mathbb{R}^n}(0,x)^2)).$$

By assuming $\tau > 0$ is sufficiently small, we see that $f(x) = d_{\mathbb{R}^n}(x, p)^2$ near $\partial B_{\tau}(p)$ and $f(x) = d_{\mathbb{R}^n}(x, 0)^2$ near $\partial B_1(0)$. Note that for any point $q \in \mathbb{R}^n$,

$$\nabla^{\Sigma}(d_{\mathbb{R}^n}(x,q)^2) = 2((x-q) - \langle x-q,N \rangle N)$$
$$(D_{\Sigma}^2(d_{\mathbb{R}^n}(x,q)^2))_x(v,v) = 2\left(|v|^2 - \operatorname{II}_{\Sigma}(x)(v,v) \langle x-q,N \rangle\right)$$

where N is any choice of normal vector at x and v is any vector in $T_x\Sigma$. Thus, we compute

$$(D_{\Sigma}^{2}f)_{x}(v,v) = 2\chi \left(|v|^{2} - \mathrm{II}_{\Sigma}(x)(v,v) \langle x - p, N \rangle \right) + 2(1-\chi) \left(|v|^{2} - \mathrm{II}_{\Sigma}(x)(v,v) \langle x, N \rangle \right) + 2\chi' (d_{\mathbb{R}^{n}}(x,p)^{2} - d_{\mathbb{R}^{n}}(x,0)^{2}) \left(|v|^{2} - \mathrm{II}_{\Sigma}(x)(v,v) \langle x, N \rangle \right)$$

$$+ 4\chi''(d_{\mathbb{R}^{n}}(x,p)^{2} - d_{\mathbb{R}^{n}}(x,0)^{2})(\langle x,v \rangle - \langle x,N \rangle \langle N,v \rangle)^{2} + 8\chi'(\langle x-p,v \rangle - \langle x-p,N \rangle \langle N,v \rangle)(\langle x,v \rangle - \langle x,N \rangle \langle N,v \rangle) - 8\chi'(\langle x,v \rangle - \langle x,N \rangle \langle N,v \rangle)^{2} = 2(|v|^{2} - \Pi_{\Sigma}(x)(v,v) \langle x-p,N \rangle) - 2(1-\chi)\Pi_{\Sigma}(x)(v,v) \langle p,N \rangle + 2\chi'(d_{\mathbb{R}^{n}}(x,p)^{2} - d_{\mathbb{R}^{n}}(x,0)^{2})(|v|^{2} - \Pi_{\Sigma}(x)(v,v) \langle x,N \rangle) + 4\chi''(d_{\mathbb{R}^{n}}(x,p)^{2} - d_{\mathbb{R}^{n}}(x,0)^{2})(\langle x,v \rangle - \langle x,N \rangle \langle N,v \rangle)^{2} - 8\chi'(\langle p,v \rangle - \langle p,N \rangle \langle N,v \rangle)(\langle x,v \rangle - \langle x,N \rangle \langle N,v \rangle).$$

Observe that for $\tau > 0$ sufficiently small, $|II_{\Sigma}|(x)| \leq \frac{1}{2}$ on the supports of $1 - \chi$, χ' and χ'' . In particular, it is easy to see that on $\Sigma \setminus B_{\tau}(p)$,

$$(D_{\Sigma}^2 f)_x(v,v) \ge 2\left(|v|^2 - \operatorname{II}_{\Sigma}(x)(v,v) \langle x-p,N \rangle\right) - Cd_{\mathbb{R}^3}(p,0)|v|^2,$$

for some C > 0 independent of τ . Combined with the assumed second fundamental form bounds, we have that

$$(D_{\Sigma}^2 f)_x(v,v) \ge 2\left(\frac{3}{4} - Cd_{\mathbb{R}^3}(p,0)\right)|v|^2.$$

Thus, as long as $d_{\mathbb{R}^3}(p,0) \leq \tau$ is sufficiently small, this is strictly positive.

Choosing such a τ , any critical point of f in $\Sigma \setminus B_{\tau}(p)$ must be a strict local minimum. The mountain pass lemma then implies that f cannot have any critical points in the interior of $\Sigma \setminus B_{\tau}(p)$. Thus, the result follows from standard Morse theory.

4. Degeneration of bounded index minimal surfaces in three-manifolds

Let I be a natural number. In this section, we analyze how a sequence of embedded minimal surfaces with index at most I in a three-manifold might degenerate and prove Theorem 1.17. By the curvature estimates from Corollary 2.3, we will be mostly working on small scales near a finite set of at most I points so that we will frequently find ourselves in situations where the following hypothesis, which we will call (\aleph) , hold.

Suppose that g_j is a sequence of metrics on $\{|x| \leq 2r_j\} \subset \mathbb{R}^3$ with $r_j \to \infty$, so that g_j is locally smoothly converging to the Euclidean metric $g_{\mathbb{R}^3}$. Assume also that:

- (1) We have $\Sigma_j \subset B_{r_j}(0)$ a sequence of properly embedded minimal surfaces with $\partial \Sigma_j \subset \partial B_{r_j}(0)$.
- (2) The surfaces have $index(\Sigma_i) \leq I$.
- (3) There is a sequence of non-empty smooth blow-up sets $\mathcal{B}_j \subset B_{\tau_0}(0)$ (where τ_0 is fixed in Lemma 3.1) with $|\mathcal{B}_j| \leq I$ and C > 0 so that

$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,\mathcal{B}_j\cup\partial\Sigma_j)\leq C,$$

for $x \in \Sigma_j$.

(4) The smooth blow-up sets converge to a set of points \mathcal{B}_{∞} and there is a smooth lamination $\Lambda \subset \mathbb{R}^3 \setminus \mathcal{B}_{\infty}$ so that Σ_j converges locally smoothly to Λ away from \mathcal{B}_{∞} .

Then, we will say that Σ_i satisfies (\aleph).

All the statements we will prove when working under hypothesis (\aleph) will turn out to be open conditions, so the reader may think of all the metric balls to be defined using the Euclidean distance.

It is convenient to write Σ'_j for the union of components of $\Sigma_j \cap B_2(0)$ which contain at least one point in \mathcal{B}_j , and Σ''_j for the union of components which contain no points in \mathcal{B}_j . Whenever j is sufficiently large, these will represent respectively the neck and the disk components of Σ_j respectively.

Lemma 4.1. The surfaces Σ''_i have uniformly bounded curvature, i.e.

$$\limsup_{j \to \infty} \sup_{x \in \Sigma_j''} |\mathrm{II}_{\Sigma_j}|(x) < \infty.$$

Proof. After passing to a subsequence, suppose that $p_j \in \Sigma_j''$ satisfies

$$|\mathrm{II}_{\Sigma_j}|(z_j) = \sup_{x \in \Sigma_j''} |\mathrm{II}_{\Sigma_j}|(x) := \lambda_j'' \to \infty.$$

Then,

$$\overline{\Sigma_j''} := \lambda_j''(\Sigma_j'' - z_j)$$

will converge to a complete, non-flat, properly embedded two-sided minimal surface $\overline{\Sigma_{\infty}''}$ in \mathbb{R}^3 with finite index (cf. Theorem B.1). On the other hand, we claim that after passing to a subsequence,

$$\overline{\Sigma'_j} := \lambda''_j (\Sigma'_j - z_j)$$

converges away from some finite set of points $\overline{\mathcal{B}}_{\infty}$ to a non-empty smooth lamination $\overline{\Lambda'}$ of $\mathbb{R}^3 \setminus \overline{\mathcal{B}}_{\infty}$. The reason that $\overline{\Lambda'}$ is non-empty is that

$$\limsup_{j \to \infty} \min_{p \in \mathcal{B}_j} \lambda_j'' d_{g_j}(z_j, p) < \infty$$

by the curvature estimates assumed in (\aleph). Thus, at least one point in the rescaled blow-up sets must remain at a bounded distance from the origin. Corollary C.5 and then Theorem B.1 imply that $\overline{\Lambda'}$ contains either a plane or properly embedded minimal surface with finite total curvature.

Because $\overline{\Sigma_{\infty}''}$ is non-flat, this is a contradiction (cf. Corollary B.2).

We give an example to illustrate the behavior described in the following two propositions in Appendix D.

Proposition 4.2 (One point of curvature concentration). Suppose that Σ_j satisfies (\aleph) with $|\mathcal{B}_j| = 1$ for each j. Then, the lamination Λ extends across \mathcal{B}_{∞} to a smooth lamination $\widetilde{\Lambda} \subset \mathbb{R}^3$. Moreover, for j sufficiently large:

(1) The surfaces Σ''_i are minimal disks of uniformly bounded curvature.

- (2) The surfaces Σ'_i intersect $\partial B_1(0)$ transversely in at most $\frac{3}{2}(I+1)$ circles.
- (3) The surfaces Σ'_i have genus at most $\frac{3}{2}(I+1)$.
- (4) The surfaces Σ'_i have uniformly bounded area, i.e.,

$$\limsup_{j \to \infty} \operatorname{area}(\Sigma'_j) < \infty.$$

(5) The surfaces Σ'_i are two-sided.

After a rotation, $\widetilde{\Lambda} = \mathbb{R}^2 \times K$ for $K \subset \mathbb{R}$ closed and $\Sigma'_j \cap B_1(0)$ smoothly converges away from \mathcal{B}_{∞} to $B_1(0) \cap (\mathbb{R}^2 \times \{\eta\})$ with finite multiplicity, for some $|\eta| \leq \frac{1}{2}$.

Proof. Let us write $\mathcal{B}_j = \{p_j\}$, $\mathcal{B}_{\infty} = \{p_{\infty}\}$ and $\lambda_j := |II_{\Sigma_j}|(p_j)$. By the definition of a smooth blow-up set, after passing to a subsequence, the surfaces

$$\overline{\Sigma}_j := \lambda_j (\Sigma_j - p_j)$$

converge to $\overline{\Sigma}_{\infty} \subset \mathbb{R}^3$, a complete, non-flat, properly embedded (and thus two-sided) minimal surface. It has index at most *I*. By⁹ [CM14], the genus *g* and number of ends *r* of $\overline{\Sigma}_j$ are both bounded by $\frac{3}{2}(I+1)$. Choose R > 0 so that $\overline{\Sigma}_{\infty}$ intersects $\partial B_R(0)$ transversely and

(4.1)
$$|\mathrm{II}_{\overline{\Sigma}_{\infty}}|(x)d_{\mathbb{R}^3}(x,0) < \frac{1}{4}$$

for $x \in \overline{\Sigma}_{\infty} \setminus B_R(0)$.

First, assume that there is $\delta > 0$ so that for j sufficiently large,

(4.2)
$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,p_j) < \frac{1}{4}$$

for $x \in \Sigma_j \cap (B_{\delta}(p_j) \setminus B_{R/\lambda_j}(p_j))$. By Lemma 3.1, we see that for j sufficiently large, the surface $\Sigma'_j \cap (B_{\delta}(p_j) \setminus B_{R/\lambda_j}(p_j))$ is topologically the union of r annuli. In particular, because $\overline{\Sigma}_{\infty}$ is two-sided, so is $\Sigma'_j \cap B_{\delta}(p_j)$.

Passing to a subsequence, let $\Lambda' \subset (B_{\delta}(p_{\infty}) \setminus \{p_{\infty}\})$ denote the lamination limit of $\Sigma'_{j} \cap B_{\delta}(p_{\infty})$ away from p_{∞} . Because $\Sigma'_{j} \cap B_{\delta}(p_{\infty})$ is two-sided and has index at most I, we claim there exists $\varepsilon \in (0, \delta]$ so that each leaf in $\Lambda' \cap B_{\varepsilon}(p_{\infty})$ has stable universal cover: if a leaf of $\Lambda' \cap B_{\delta}(p_{\infty})$ has the convergence to occurring with multiplicity bigger than one, then it must have stable universal cover (cf. [MR06, Lemma A.1]). On the other hand, if we consider a leaf $\Lambda' \cap B_{\delta}(p_{\infty})$ where the convergence to occurs with multiplicity one, then this leaf itself must have bounded index. Moreover, the sum of the index of such leaves must be bounded above by I, or else it would violate the bound for $\operatorname{index}(\Sigma'_{j})$, and in particular there are only finitely many unstable leaves. Since for each leaf we may argue as in [FC85, Proposition 1] to find $\varepsilon \in (0, \delta]$ so that it is stable in $B_{\varepsilon}(p_{\infty})$ and there are only finitely many of such leaves, the claim follows. Thus, by Proposition C.3, $\Lambda' \cap B_{\varepsilon}(p_{\infty})$ extends across p_{∞} .

By Lemma 4.1, we see that $\Sigma_j'' \cap B_{\varepsilon}(p_{\infty})$ has a smooth lamination limit. Putting these two facts together, we see that the limit lamination of Σ_j away from p_{∞} , say $\Lambda \subset \mathbb{R}^3 \setminus \{p_{\infty}\}$, must extend across p_{∞} to a smooth lamination $\widetilde{\Lambda} \subset \mathbb{R}^3$. Arguing as above, we see that $\widetilde{\Lambda}$ has finite index. Thus, by Corollary B.2, $\widetilde{\Lambda}$ is either a non-flat

18

⁹We remark that it is not strictly necessary to refer to [CM14] here. Indeed, one could argue in a similar manner to the proof of Theorem 1.2 to use our blow-up strategy along with the fact [FC85] that "a finite index surface in \mathbb{R}^3 cannot have infinite genus" to prove that there is C = C(I) so that an embedded minimal surface in \mathbb{R}^3 with index at most I has at most C(I) genus and ends. Referring to [CM14] allows us to avoid such a discussion (and also allows us to obtain functions m(I) and r(I) in Theorem 1.17 that are explicitly computable).

single properly embedded surface of finite total curvature, or else is a lamination of \mathbb{R}^3 by parallel planes.

In the first case, the convergence of Σ_j to $\widetilde{\Lambda}$ must occur with multiplicity one (otherwise $\widetilde{\Lambda}$ would be stable and thus a plane; see e.g., [MR06, Lemma A.1]). In this case, Allard's regularity theorem [All72] would imply that Σ_j converged smoothly to $\widetilde{\Lambda}$ near p_{∞} . This is not compatible with the fact that the curvature of Σ_j at p_j is blowing up, so we see that $\widetilde{\Lambda}$ is a (non-empty) lamination of \mathbb{R}^3 by parallel planes. In particular, (1) follows from this and Lemma 4.1.

We will rotate the coordinates so that $\widetilde{\Lambda} = \mathbb{R}^2 \times K$ for $K \subset \mathbb{R}$ a non-empty closed set. Hence, we see that for j large, (4.2) actually holds for all $x \in \Sigma_j \cap (B_2(0) \setminus B_{R/\lambda_j}(p_j))$. From this, assertions (2) through (5) follow easily from Lemma 3.1 (note that we have assumed that $\mathcal{B}_j \subset B_{\tau_0}(0)$ and $\tau_0 < \frac{1}{2}$) and the above description of $\overline{\Sigma}_{\infty}$.

Now, it remains to show that we can find $\delta > 0$ so that (4.2) holds for $x \in \Sigma_j \cap (B_{\delta}(p_j) \setminus B_{R/\lambda_j}(p_j))$. If this were to fail, for j large, we can find $\delta_j \to 0$ so that $\delta_j > R/\lambda_j$ and (4.2) holds for all $x \in \Sigma_j \cap (B_{\delta_j}(p_j) \setminus B_{R/\lambda_j}(p_j))$, but δ_j is the largest possible number with this property. Observe that this choice of δ_j guarantees that

(4.3)
$$\sup_{z \in \partial B_{\delta_j}(p_j) \cap \Sigma_j} |\Pi_{\Sigma_j}|(z) d_{g_j}(z, p_j) = \frac{1}{4}.$$

We now consider

$$\hat{\Sigma}_j = \delta_j^{-1} (\Sigma_j - p_j).$$

Note that the curvature of $\hat{\Sigma}_j$ at the origin cannot be uniformly bounded as $j \to \infty$, as otherwise $\hat{\Sigma}_j$ would limit to a rescaled version of $\overline{\Sigma}_{\infty}$. This would contradict the choice of R and δ_j , in particular (4.1). Hence, $\hat{\Sigma}_j$ satisfies all of the hypothesis of the proposition, in addition to satisfying (the rescaled version of) (4.2) on $B_1(0)$, i.e.

$$|\mathrm{II}_{\hat{\Sigma}_j}|(x)d_{\hat{g}_j}(x,0) < \frac{1}{4}$$

for $x \in B_1(0) \cap \hat{\Sigma}_j$. By the above argument, $\hat{\Sigma}_j$ converges subsequentially to a lamination of \mathbb{R}^3 by parallel planes. The (scale invariant) curvature estimates assumed in (\aleph) and the assumption that $|\mathcal{B}_j| = 0$ guarantee that

$$II_{\hat{\Sigma}_{i}}|(x)d_{\hat{g}_{j}}(x,0) \le C$$

for, e.g., $x \in B_2(0) \cap \hat{\Sigma}_j$. Thus, the convergence of $\hat{\Sigma}_j$ to the lamination by parallel planes takes place smoothly away from $\{0\}$. In particular,

$$\sup_{\epsilon \partial B_1(0) \cap \hat{\Sigma}_j} |\mathrm{II}_{\hat{\Sigma}_j}|(z) \to 0.$$

This contradicts (4.3), after rescaling.

Proposition 4.3 (Multiple points of curvature concentration). There are functions m(I) and r(I) so that the following holds. Suppose that Σ_j satisfies (\aleph). Then, the lamination Λ extends across \mathcal{B}_{∞} to a smooth lamination $\widetilde{\Lambda} \subset \mathbb{R}^3$. Moreover, for j sufficiently large:

(1) The surfaces Σ''_i are minimal disks of uniformly bounded curvature.

- (2) The surfaces Σ'_i intersect $\partial B_1(0)$ transversely in at most m(I) circles.
- (3) The surfaces Σ'_i have genus at most r(I).
- (4) The surfaces Σ'_i have uniformly bounded area, i.e.,

$$\limsup_{j \to \infty} \operatorname{area}(\Sigma'_j) < \infty$$

(5) The surfaces Σ'_{j} are two-sided.

After a rotation, $\widetilde{\Lambda} = \mathbb{R}^2 \times K$ for $K \subset \mathbb{R}$ closed and $\Sigma'_j \cap B_1(0)$ converges to $B_1(0) \cap (\mathbb{R}^2 \times \{\eta_1, \ldots, \eta_n\})$ with finite multiplicity for some $|\eta_i| \leq \frac{1}{2}$.

Proof. We will induct on the index bound I in (\aleph). If I = 1, then the proposition follows from Proposition 4.2 above. Now, assume that the proposition holds for I - 1 and that $index(\Sigma_i) \leq I$.

We first consider the case that $|\mathcal{B}_{\infty}| \geq 2$. Pick $\delta > 0$ so that \mathcal{B}_{∞} is 4 δ -separated. In particular, $B_{\delta}(\mathcal{B}_{\infty})$ is a disjoint union of balls and, after passing to a subsequence, we may assume that for any connected component B of $B_{\delta}(\mathcal{B}_{\infty})$

$$\operatorname{index}(\Sigma_j \cap B) \ge 1.$$

Because we are assuming that $|\mathcal{B}_{\infty}| \geq 2$, this implies that

$$\operatorname{index}(\Sigma_j \cap B) \le I - 1.$$

Now, we choose $\varepsilon_j \to 0$ sufficiently slowly so that $\mathcal{B}_j \subset B_{\varepsilon_j \tau_0/2}(\mathcal{B}_\infty)$ and

$$\liminf_{j \to \infty} \varepsilon_j \min_{p \in \mathcal{B}_j} |\mathrm{II}_{\Sigma_j}|(p) = \infty$$

We claim that (after taking $\delta > 0$ smaller if necessary) for j large we have

(4.4)
$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,\mathcal{B}_{\infty}) < \frac{1}{4}$$

for $x \in \Sigma_j \cap (B_{\delta}(\mathcal{B}_{\infty}) \setminus B_{\varepsilon_j}(\mathcal{B}_{\infty}))$. If this were to fail, then we may argue as in the one point case, Proposition 4.2: choose some component $B = B_{\delta}(p_{\infty})$ of $B_{\delta}(\mathcal{B}_{\infty})$ (for some $p_{\infty} \in \mathcal{B}_{\infty}$), where it failed, and choose the largest $\delta_j \geq \varepsilon_j$ so that (4.4) holds for $x \in \Sigma_j \cap (B_{\delta_j}(p_{\infty}) \setminus B_{\varepsilon_j}(p_{\infty}))$. Thus, the surfaces

$$\delta_j^{-1}(\Sigma_j \cap B_\delta(p_\infty) - p_\infty)$$

satisfy the inductive hypothesis and so, after passing to a subsequence, they converge to a lamination of \mathbb{R}^3 by parallel planes, contradicting the choice of δ_j .

Because we now know that (4.4) holds, we are able to transfer topological information from the scale $B_{\varepsilon_j}(\mathcal{B}_{\infty})$ (where we may apply the inductive hypothesis, by choice of ε_j) to $B_{\delta}(\mathcal{B}_{\infty})$, using Lemma 3.1. In particular, we see that any component of $\Sigma_j \cap B_{\delta}(\mathcal{B}_{\infty})$ containing some point in \mathcal{B}_j intersects $B_{\delta}(\mathcal{B}_{\infty})$ transversely in at most m(I-1) circles, has genus at most r(I-1) and is is two-sided. Thus, as in the proof of Proposition 4.2, we may use Proposition C.3 to see that Λ has removable singularities, i.e. $\Lambda = \tilde{\Lambda} \setminus \mathcal{B}_{\infty}$ for a smooth lamination $\tilde{\Lambda} \subset \mathbb{R}^3$ with finite index. By Corollary B.2, we see that $\tilde{\Lambda}$ is a lamination by parallel planes. In particular, assertion (1) follows from this and Lemma 4.1. Moreover, it is not hard to check that $\Sigma'_i \cap (B_1(0) \setminus B_{\delta}(\mathcal{B}_{\infty}))$ satisfies the hypothesis of Lemma A.3, which yields the asserted bounds on the genus and number of boundary components of Σ'_j . Combined with the structure of $\tilde{\Lambda}$, this also yields the asserted area bounds, so we have that assertions (2)-(5) also hold.

Thus, it remains to consider the case that $|\mathcal{B}_{\infty}| = 1$. Passing to a subsequence, we may assume that $|\mathcal{B}_j| \geq 2$ for each j, as otherwise we could apply Proposition 4.2. Then, we may choose $p_j, q_j \in \mathcal{B}_j$ so that

$$\varepsilon_j \tau_0/2 := d_{g_j}(p_j, q_j) = \max_{\substack{p, q \in \mathcal{B}_j \\ p \neq q}} d_{g_j}(p, q) \to 0.$$

Then, consider the sequence

$$\overline{\Sigma}_j := \varepsilon_j^{-1} (\Sigma_j - p_j)$$

By definition of a sequence of smooth blow-up sets (i.e., the various points cannot appear in the blow-up of the other points), the curvature must still be blowing up at each point in $\overline{\mathcal{B}}_j = \varepsilon_j^{-1}(\mathcal{B}_j - p_j)$. Thus $\overline{\Sigma}_j$ satisfies the hypothesis of the proposition with $|\overline{\mathcal{B}}_{\infty}| \geq 2$, so the conclusion of the proposition holds for $\overline{\Sigma}_j$. At this point, we may argue as above (cf. the analogous argument in the proof of Proposition 4.2), establishing the curvature estimate (4.4) for $x \in \Sigma_j \cap (B_{\delta}(p_j) \setminus B_{\varepsilon_j}(p_j))$ for some $\delta > 0$. As before, this allows us to remove the singularities in the limit lamination and conclude that it must be a lamination by planes. Using this, we may readily transfer the topological information out to the scale of $B_1(0)$ for Σ_j using Lemma 3.1 and as before conclude assertions (1) through (5).

4.1. Completing the proof of Theorem 1.17. Assume that $\Sigma_j \subset (M^3, g)$ is a sequence of closed embedded minimal surfaces with

$$\operatorname{index}(\Sigma_j) \leq I.$$

Passing to a subsequence, Corollary 2.3 yields a sequence of smooth blow-up sets \mathcal{B}_j so that $|\mathcal{B}_j| \leq I$ and a constant C > 0 so that

$$|\mathrm{II}_{\Sigma_j}|(x)\min\{1, d_g(x, \mathcal{B}_j)\} \le C$$

Passing to a further subsequence, \mathcal{B}_j converges to a finite set of points \mathcal{B}_{∞} and Σ_j converges away from \mathcal{B}_{∞} to a lamination $\mathcal{L} \subset M \setminus \mathcal{B}_{\infty}$. The remaining argument is very similar to the proof of Proposition 4.3, so we omit some of the details below.

Choose $\varepsilon_j \to 0$ sufficiently slowly so that $\mathcal{B}_j \subset B_{\varepsilon_j \tau_0/2}(\mathcal{B}_\infty)$ and

$$\liminf_{j \to \infty} \varepsilon_j \min_{p \in \mathcal{B}_j} |\mathrm{II}_{\Sigma_j}|(p) = \infty.$$

We claim that by taking $\varepsilon_0 > 0$ sufficiently small (in particular, so that it is smaller than the injectivity radius and so that \mathcal{B}_{∞} is $4\varepsilon_0$ -separated), for j large, we have the improved curvature bounds

(4.5)
$$|\mathrm{II}_{\Sigma_j}|(x)d_g(x,\mathcal{B}_{\infty}) < \frac{1}{4}$$

for $x \in \Sigma_j \cap (B_{2\varepsilon_0}(\mathcal{B}_\infty) \setminus B_{\varepsilon_j}(\mathcal{B}_\infty))$. If not, we may pick a connected component $\Sigma_j \cap (B_{2\varepsilon_0}(\mathcal{B}_\infty) \setminus B_{\varepsilon_j}(\mathcal{B}_\infty))$ where it fails and rescale the largest ball in which (4.5) is satisfied to unit scale. This rescaled sequence then satisfies the hypothesis of Proposition 4.3, so we see that it limits to a lamination of \mathbb{R}^3 by parallel planes (away from a discrete set).

This contradicts the fact that we chose the largest ball satisfying (4.5). Thus, we may find some $\varepsilon_0 > 0$ as claimed. Taking $\varepsilon_0 > 0$ even smaller if necessary, we may arrange that for every component *B* of $B_{2\varepsilon_0}(\mathcal{B}_{\infty})$, the metric *g* restricted to *B* and rescaled by by ε_0^{-1} around its center satisfies the hypothesis in Lemma 3.1.

Now, Propositions 4.2 and 4.3 applied to each component B of $\Sigma_j \cap B_{\varepsilon_j}(\mathcal{B}_{\infty})$, after rescaling it by ε_j^{-1} around the center of B yields the desired topological information at the scale of $B_{\varepsilon_j}(\mathcal{B}_{\infty})$. The improved curvature estimates in (4.5) and Lemma 3.1 then allow us to transfer this information out to the scale of $B_{\varepsilon_0}(\mathcal{B}_{\infty})$, exactly as in the proof of Propositions 4.2 and 4.3. In particular, topological statements (1.a), (1.b), (1.c) follow, and also (2.a) by Lemma 4.1. Then, each component of $\Sigma_j \cap B_{\varepsilon_0}(\mathcal{B}_{\infty})$ either has uniformly bounded curvature, or has uniformly bounded area and is twosided, so we may apply Proposition C.1 to conclude that the lamination \mathcal{L} extends to a smooth lamination $\widetilde{\mathcal{L}}$ with $\mathcal{L} = \widetilde{\mathcal{L}} \setminus \mathcal{B}_{\infty}$.

Finally, fix $\varepsilon \in (0, \varepsilon_0]$ and $B = B_{\varepsilon_0}(p_{\infty})$ a connected component component of $B_{\varepsilon_0}(\mathcal{B}_{\infty})$. Because $\widetilde{\mathcal{L}}$ is smooth in B, as $\varepsilon \to 0$, each leaf in $\varepsilon^{-1}(\widetilde{\mathcal{L}} \cap B - p_{\infty})$ converges with multiplicity one to a plane in \mathbb{R}^3 . Rotating a local coordinate frame, we may assume that all such planes are of the form $\mathbb{R}^2 \times \{t\}$ for some $t \in \mathbb{R}$.

Thus, by (1.a) we can see that for j sufficiently large (depending on ε) any component of $\Sigma_j \cap B_{2\varepsilon}(p_{\infty})$ must intersect $B_{\varepsilon}(p_{\infty}) \setminus B_{\varepsilon^2}(p_{\infty})$ union of at most m(I) annuli, which converge graphically to the annulus $(\mathbb{R}^2 \times \{0\}) \cap (B_{\varepsilon}(p_{\infty}) \setminus B_{\varepsilon^2}(p_{\infty}))$. Combined with the monotonicity formula, the area estimate (1.d) easily follows. The argument for (2.b) follows a similar line of reasoning, except any neck region is converging smoothly everywhere to a leaf in $\widetilde{\mathcal{L}} \cap B$, which is nearly planar on small scales.

5. SURGERY FOR BOUNDED INDEX SURFACES IN THREE-MANIFOLDS

In this section, we describe how Corollary 1.19 follows from Theorem 1.17. We first prove the following local description of the surgery operation.

Proposition 5.1 (Local picture of surgery). Suppose that Γ_j is a sequence of embedded surfaces in $B_3(0)$ with $\partial \Gamma_j \subset \partial B_3(0)$, and so that:

- (1) The surfaces $\Gamma_j \setminus \overline{B_1(0)}$ converge smoothly, with finite multiplicity, to the flat annulus $A(3,1) := \left(B_3(0) \setminus \overline{B_1(0)}\right) \cap \{x^3 = 0\}$ as $j \to \infty$.
- (2) The set of components of Γ_j which are topological disks converge smoothly to the flat disk $D(3) := B_3(0) \cap \{x^3 = 0\}$ as $j \to \infty$.

Then, for j sufficiently large, we may construct embedded surfaces $\widetilde{\Gamma}_j$ with $\partial \widetilde{\Gamma}_j \subset \partial B_3(0)$, and so that:

- (1) The surfaces Γ_i agree with the Γ_i near $\partial B_3(0)$.
- (2) Any component of Γ_j which is topologically a disk is unchanged.
- (3) The surfaces $\widetilde{\Gamma}_j$ converge smoothly, with finite multiplicity, to the flat disk D(3) as $j \to \infty$.

Proof. Fix a smooth cutoff function $\chi : \mathbb{R}^2 \to [0,1]$, with $\chi(x) \equiv 1$ for $|x| \ge \frac{7}{4}$ and $\chi(x) \equiv 0$ for $|x| \le \frac{5}{4}$.

We define the cylinder and annular cylinder

$$C(r) := \{ (x^1, x^2, x^3 : (x^1)^2 + (x^2)^2 < r^2 \}, \quad r > 0,$$

$$C(r_1, r_2) := C(r_1) \setminus \overline{C(r_2)}, \quad r_1 > r_2 > 0.$$

Taking j sufficiently large, each component of $\Gamma_j \cap C(2,1)$ is graphical over the flat annulus A(2,1), and the topological disk components of $\Gamma_j \cap C(2)$ are graphical over the flat disk D(2).

For now, we assume at most two of the components of Γ_j are topological disks, and each of the disk components, if they exist, is either the topmost component or bottommost component. Choose a smooth function $w_j: D(2) \to \mathbb{R}$ so that

- (1) The graph of w_i is contained in $B_3(0)$.
- (2) The graph of w_j lies strictly above (resp. below) the bottommost (resp. uppermost) disk if it exists.
- (3) The function w_j converges smoothly to 0 as $j \to \infty$.

For example, when Γ_j contains both an uppermost and bottommost disk, then we may take the average of their respective graphs. We additionally choose real numbers $\eta_j \to 0$ so that the graph of $w_j + \eta$ satisfies the above properties as well for all $\eta \in (0, \eta_j)$.

We may find functions $u_{j,1}, \ldots, u_{j,n(j)} : A(2,1) \to \mathbb{R}$ so that any non-disk component of Γ_j is the graph of the $u_{j,l}$ in C(2,1). By assumption, for all k,

$$\sup_{\in\{1,\dots,n(j)\}} \|u_{j,l}\|_{C^k(A(2,1))} \to 0$$

 $l \in \{1, ..., n(j)\}$ as $j \to \infty$. By embeddedness of Γ_j , we may arrange that

$$u_{j,1}(x) < u_{j,2}(x) < \dots < u_{j,n(j)}(x)$$

for $x \in A(2, 1)$.

Now, we define

$$\widetilde{u}_{j,l}(x) = \chi(x)u_{j,l}(x) + (1 - \chi(x))\left(w_j(x) + \frac{l}{n(j)}\eta_j\right)$$

We now define a surface $\widetilde{\Gamma}_j$ which agrees with Γ_j in $B_3(0) \setminus C(2)$ and which is defined inside of C(2) to be the union of the graphs of the $\widetilde{u}_{j,l}$ along with the disk components in Γ_j , if they exist. It is easy to check that $\widetilde{\Gamma}_j$ satisfies the asserted properties.

Finally, we may easily reduce the case of general Γ_j to the above case by considering contiguous subsets of the components of Γ_j which are in the above form and applying the argument above to the maximal such subsets. This choice will preserve embeddedness, because we have chosen them so that there will at least be a disk separating the non-disk components of different subsets.

Now, we may complete the proof of the surgery result. Consider Σ_j a sequence of compact embedded minimal surfaces in (M^3, g) with $index(\Sigma_j) \leq I$. We pass to a subsequence so that the conclusion of Theorem 1.17 applies. In particular, there is a

finite set of points $\mathcal{B}_{\infty} \subset M$ with $|\mathcal{B}_{\infty}| \leq I$ and a smooth lamination $\widetilde{\mathcal{L}}$ of M so that Σ_i converges to $\mathcal{L} = \widetilde{\mathcal{L}} \setminus \mathcal{B}_{\infty}$ away from \mathcal{B}_{∞} .

Take ε_0 as in Theorem 1.17 and choose $\varepsilon \in (0, \varepsilon_0]$. Pick any $p \in \mathcal{B}_{\infty}$; we will show how to perform the surgery in $B_{\varepsilon}(p)$. Write L for the leaf of $\widetilde{\mathcal{L}} \cap B_{\varepsilon}(p)$ that passes through p. We may fix a diffeomorphism of $\Psi : B_{\varepsilon}(p) \to B_3(0) \subset \mathbb{R}^3$ so that Ψ maps $B_{\varepsilon/3}(0)$ diffeomorphically onto $B_1(0)$ and L onto the flat disk $D(3) \subset \mathbb{R}^3$ as in Proposition 5.1.

Consider the connected components of $\Sigma_j \cap B_{\varepsilon}(p_i)$ which are converging smoothly to L in the annulus $B_{\varepsilon}(p) \setminus \overline{B_{\varepsilon/3}(p)}$ (by Theorem 1.17, this includes all of the neck components, i.e., all of the components of $\Sigma_j \cap B_{\varepsilon}(p)$ containing some point in \mathcal{B}_j). Using the maximum principle, the area bounds and curvature estimates for the disk components show that they converge smoothly to L (although they might do so with infinite multiplicity). Now, we define Γ_j to be the union of all of the neck components of $\Sigma_j \cap B_{\varepsilon}(p)$, as well as all of the disc components which are directly adjacent (either above or below) to a neck component.

It is not hard to see that if the uppermost (resp. lowermost) component of Γ_j is a neck component, we may simply add in a disk which is above (resp. below) all of the components of Γ_j , but which is below (resp. above) all of the disk components not converging to L.

Now we apply Proposition 5.1 to Γ_j (and then removing the extra disks on top and bottom, if we had to add them) and replace $\Sigma_j \cap B_{\varepsilon}(p)$ by the resulting surface. Repeating this for each $p \in \mathcal{B}_{\infty}$ yields $\widetilde{\Sigma}_j$. The asserted properties of $\widetilde{\Sigma}_j$ follow easily from Proposition 5.1 and Theorem 1.17.

6. PROOFS OF THE THREE-DIMENSIONAL COMPACTNESS RESULTS

Proof of Theorem 1.1 for n = 3. Fix $I \in \mathbb{N}$, $A < \infty$, and a closed Riemannian threemanifold (M,g). Suppose that $\Sigma_j \subset (M,g)$ is a sequence of connected, embedded, closed minimal surfaces with $\operatorname{index}(\Sigma_j) \leq I$ and $\operatorname{area}(\Sigma_j) \leq A$ but $\operatorname{genus}(\Sigma_j) \to \infty$. By¹⁰ Corollary 1.19, we may find $\widetilde{\Sigma}_j$ with uniformly bounded area and curvature, but so that

$$\operatorname{genus}(\Sigma_j) \ge \operatorname{genus}(\Sigma_j) - \tilde{r}(I) \to \infty.$$

This is a contradiction: after passing to a subsequence, the surfaces Σ_j must converge smoothly and with finite multiplicity to some closed, embedded minimal surface $\widetilde{\Sigma}_{\infty}$ (which must have finite genus).

Proof Theorem 1.3. Fix $I \in \mathbb{N}$ and (M, g) a closed Riemannian three-manifold with positive scalar curvature. We only need to prove the area bound, since the genus bound would immediately follow from Theorem 1.1 (the case n = 3 is proven above). Suppose that $\Sigma_j \subset (M, g)$ is a sequence of connected, closed, embedded minimal surfaces with index $(\Sigma_j) \leq I$ and area $_g(\Sigma_j) \to \infty$.

¹⁰Note that the full statement Corollary 1.19 is not really needed to prove this, one may directly use the description from Theorem 1.17 to prove the claim, somewhat like Theorems 1.1 or 1.2. However, the argument using Corollary 1.19 is very short, which is why we have presented the proof in this form.

After passing to a subsequence, by Theorem 1.17, there is a finite set of points \mathcal{B}_{∞} and a lamination $\widetilde{\mathcal{L}} \subset M$ so that Σ_j converges locally to the lamination $\mathcal{L} := \widetilde{\mathcal{L}} \setminus \mathcal{B}_{\infty}$ away from \mathcal{B}_{∞} . Because the area of Σ_j is diverging, passing to a further subsequence, there is $p \in M \setminus \mathcal{B}_{\infty}$ so that

$$\liminf_{j \to \infty} \operatorname{area}_g(\Sigma_j \cap B_r(p)) = \infty.$$

for all r > 0. A standard argument along the lines of [MR05, Lemma 1.1], [MR06, Lemma A.1], and [CCE15, Proposition 2.1] shows that there is a leaf $p \in L \subset \mathcal{L}$ with stable universal cover and so that for r > 0 fixed sufficiently small, $\Sigma_j \cap B_r(p)$ consists of $n(j) \to \infty$ sheets, which are all smoothly graphically converging to $L \cap B_r(p)$.

Because $\mathcal{L} = \mathcal{L} \setminus \mathcal{B}_{\infty}$ has removable singularities, there is a smooth complete minimal surface \tilde{L} so that $L = \mathcal{L} \setminus \mathcal{B}_{\infty}$. The log-cutoff trick shows that stability extends across isolated points, so \tilde{L} has stable universal cover \hat{L} . We think of \hat{L} as an immersed stable minimal surface in M. If we consider a disk $D \subset \hat{L}$ and if x is any point in the interior of D, by Schoen-Yau [SY82, SY83], the intrinsic distance to the boundary must satisfy:

$$d_D(x,\partial D) \le \frac{2\pi\sqrt{2}}{\sqrt{3\kappa_0}},$$

where $\kappa_0 > 0$ is the infimum of the scalar curvature of M. This implies that \widehat{L} must be compact, since D is arbitrary. By [FCS80, Theorem 3], \widehat{L} is a two-sphere.

We choose $\varepsilon > 0$ smaller than ε_0 from the surgery theorem and small enough so that $p \notin \mathcal{B}_{2\varepsilon}(\mathcal{B}_{\infty})$. Let $\widetilde{\Sigma}_j$ denote the surfaces resulting from a surgery at scale ε , as constructed in Corollary 1.19. Because the original surfaces Σ_j are connected, Corollary 1.19 implies that the number of components of $\widetilde{\Sigma}_j$ is uniformly bounded above, $|\pi_0(\widetilde{\Sigma}_j)| \leq m(I) + 1$.

Putting these facts together, we may find a connected component $\widehat{\Sigma}_j \subset \widetilde{\Sigma}_j$ so that $\operatorname{area}_g(\widehat{\Sigma}_j) \to \infty$ and so that $\widehat{\Sigma}_j \cap B_{\varepsilon}(p)$ is smoothly converging to $L \cap B_{\varepsilon}(p)$. The maximum principle then implies that $\widehat{\Sigma}_j$ converges locally smoothly to \widetilde{L} . In particular, the universal cover of $\widehat{\Sigma}_j$ converges in the sense of immersions to \widehat{L} , which we have seen is a topological sphere. This implies that the area of $\widehat{\Sigma}_j$ is uniformly bounded, a contradiction.

7. Bounded diffeomorphism type in higher dimensions

Here, we discuss the $4 \le n \le 7$ case of Theorems 1.1 and 1.2. Motivated by the three-dimensional case, we define the hypothesis (\beth) as follows.

Fix $4 \leq n \leq 7$ and suppose that g_j is a sequence of metrics on $\{|x| \leq 2r_j\} \subset \mathbb{R}^n$ that is locally smoothly converging to $g_{\mathbb{R}^n}$. Assume that:

- (1) We have $\Sigma_j \subset B_{r_j}(0) \subset \mathbb{R}^n$ a sequence of properly embedded minimal hypersurfaces with $\partial \Sigma_j \subset \partial B_{r_j}(0)$.
- (2) The surfaces Σ_i are connected.
- (3) The hypersurfaces have $index(\Sigma_i) \leq I$.

- (4) The hypersurfaces satisfy $\operatorname{vol}(\Sigma_j) \leq \Lambda r_i^{n-1}$.
- (5) There is a sequence of non-empty smooth blow-up sets $\mathcal{B}_j \subset B_{\tau_0}(0)$ (where τ_0 is fixed in Lemma 3.1) with $|\mathcal{B}_j| \leq I$ and C > 0 so that

$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,\mathcal{B}_j\cup\partial\Sigma_j)\leq C_j$$

for $x \in \Sigma_j$.

(6) The smooth blow-up sets converge to a set of points \mathcal{B}_{∞} and for any r > 0, the hypersurfaces $\Sigma_j \cap B_r(0)$ converge in sense of varifolds to a disk with multiplicity $k \in \mathbb{N}$, i.e.

$$[\Sigma_j \cap B_r(0)] \rightharpoonup k[\{x^n = 0\} \cap B_r(0)]$$

Then, we say that Σ_i satisfies (\beth) .

Let us briefly note that the main difference between hypothesis (\beth) and the hypothesis (\aleph) used in three dimensions is the assumption that the surfaces are connected (in addition to the assumption that they satisfy a uniform area bound). The connectedness assumption is useful to compensate for the fact that the half-space theorem fails in higher dimensions. To exploit this assumption, we will work "big to small," rather than¹¹ "inside out," when proving the crucial curvature estimates, e.g., (7.1).

Proposition 7.1. Given a sequence Σ_j satisfying (\beth) that intersect $\partial B_1(0)$ transversely, we may pass to a subsequence so that all of the $\Sigma_j \cap B_1(0)$ are diffeomorphic.

Proof. We prove this by induction on I. For I = 0 this trivially follows from the curvature estimates.

We first consider the one point of concentration, i.e. $|\mathcal{B}_j| = 1$. We write $\mathcal{B}_j = \{p_j\}$ and $\mathcal{B}_{\infty} = \{p_{\infty}\}$ and $\lambda_j = |\Pi_{\Sigma_j}|(p_j)$. By passing to a subsequence, we have that

$$\overline{\Sigma}_j := \lambda_j (\Sigma_j - p_j)$$

converges to $\overline{\Sigma}_{\infty} \subset \mathbb{R}^n$ a complete, non-flat, properly embedded minimal surface with index at most I and $\operatorname{vol}(\overline{\Sigma}_{\infty} \cap B_r(0)) \leq \Lambda r^{n-1}$ (by the monotonicity formula). Because of these properties, $\overline{\Sigma}_{\infty}$ must be "regular at infinity" in the sense that outside of a large compact set, it is the finite union of a graphs, all over the same fixed plane, of functions with nice asymptotic behavior, see [Sch83b, Tys89]. In particular, we may take R > 0 so that $\overline{\Sigma}_{\infty}$ intersects $\partial B_R(0)$ transversely and

$$|\mathrm{II}_{\overline{\Sigma}_{\infty}}|(x)d_{\mathbb{R}^n}(x,0) < \frac{1}{4}$$

for $x \in \overline{\Sigma}_{\infty} \setminus B_R(0)$.

We claim that for j sufficiently large,

(7.1)
$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,p_j) < \frac{1}{4}$$

26

¹¹In three dimensions, we worked "small to big" to guarantee that any singular points of a limit lamination at the given scale must be the limit of two-sided surfaces. This allowed us to apply Proposition C.3 to remove the singularities. Thanks to the volume estimates assumed here, we do not need to know a priori that the singularities of a limit surface are two-sided to remove its singularities, see Proposition C.6.

for $x \in \Sigma_j \cap (B_2(0) \setminus B_{R/\lambda_j}(p_j))$. If this holds, then Lemma 3.1 easily is seen to imply that for j sufficiently large, all of the hypersurfaces $\Sigma_j \cap B_1(0)$ are diffeomorphic (here, we have used the obvious fact that "regular ends" are diffeomorphic to $\mathbb{S}^{n-2} \times (0,1)$ with the standard smooth structure).

On the other hand, if (7.1) does not hold, we may choose δ_j to be the smallest¹² radius greater than R/λ_j so that

$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,p_j) < \frac{1}{4}$$

holds for $x \in \Sigma_j \cap (B_2(0) \setminus B_{\delta_j}(p_j))$. Note that for j sufficiently large, such a δ_j exists and moreover $\delta_j \to 0$. This follows from fact that Σ_j converges smoothly to $\{x^n = 0\}$ away from p_{∞} .

Define

$$\hat{\Sigma}_j := \delta_j^{-1} (\Sigma_j - p_j).$$

Passing to a subsequence, there is $\hat{\Sigma}_{\infty} \subset \mathbb{R}^n$ so that $\hat{\Sigma}_j$ converges locally smoothly with finite multiplicity to $\hat{\Sigma}_{\infty}$ away from $\{0\}$, and converges in the sense of varifolds in $B_1(0)$. Because $\hat{\Sigma}_{\infty}$ has finite index, we may apply Proposition C.6 to see that the singularity at $\{0\}$ is removable. In particular (after relabeling the hypersurface), $\hat{\Sigma}_{\infty}$ is an embedded minimal hypersurface in \mathbb{R}^n with $\operatorname{index}(\hat{\Sigma}_{\infty}) \leq I$ and $\operatorname{vol}(\hat{\Sigma}_{\infty} \cap B_r(0)) \leq \Lambda r^{n-1}$. In particular, it is regular at infinity and has finitely many components. Hence, we may choose $\gamma \geq 1$ so that $\partial B_{\gamma}(0)$ intersects each component transversely, and $\hat{\Sigma}_{\infty} \cap \partial B_{\gamma}(0)$ is the disjoint union of finitely many manifolds diffeomorphic to \mathbb{S}^{n-2} with the standard smooth structure.

By choice of δ_j , the curvature estimates (7.1) hold for $x \in \Sigma_j \cap (B_2(0) \setminus B_{\gamma\delta_j}(p_j))$. Applying Lemma 3.1, we see that $\Sigma_j \cap (B_2(0) \setminus B_{\gamma\delta_j}(p_j))$ is diffeomorphic to the union of annular regions. In particular, $\Sigma_j \cap B_{\gamma\delta_j}(p_j)$ must be connected (because we have assumed that Σ_j is connected in (\square)). From this, we see that $\hat{\Sigma}_{\infty}$ is connected. Observe that the convergence of $\hat{\Sigma}_j$ to $\hat{\Sigma}_{\infty}$ cannot be smooth at $\{0\}$ by choice of R and the assumption that $R/\lambda_j \leq \delta_j$. In particular, the convergence of $\hat{\Sigma}_j$ to $\hat{\Sigma}_{\infty}$ must occur with multiplicity at least two, so $\hat{\Sigma}_{\infty}$ is (two-sided) stable and thus a plane; note that this uses the fact that $\hat{\Sigma}_{\infty}$ is connected.¹³ The convergence of $\hat{\Sigma}_j$ to $\hat{\Sigma}_{\infty}$ occurs smoothly near $\partial B_1(0)$. This contradicts the choice of δ_j (namely that (7.1) fails at some point in $\Sigma_j \cap \partial B_{\delta_j}(p_j)$). This completes the proof in the case that $|\mathcal{B}_j| = 1$.

Now, we consider the case of $|\mathcal{B}_{\infty}| \geq 2$. Pick $\delta > 0$ so that \mathcal{B}_{∞} is 4 δ -separated. In particular, if B is a component of $B_{\delta}(\mathcal{B}_{\infty})$, then for j sufficiently large, we see that

$$\operatorname{index}(\Sigma_j \cap B) \le I - 1.$$

¹²Again, we emphasize that δ_j is not chosen in the same way here as in three dimensions, e.g., the proof of Proposition 4.2. Here, we are proving the estimate (7.1) by working from big to small scales, rather than small to big.

¹³If we did not arrange for $\hat{\Sigma}_{\infty}$ to be connected, then we could only conclude that it contained a plane through the origin but it could have other non-flat components.

We may choose $\varepsilon_j \to 0$ sufficiently slowly so that $\mathcal{B}_j \subset B_{\varepsilon_i/j}(\mathcal{B}_\infty)$,

$$\liminf_{j \to \infty} \varepsilon_j \min_{p \in \mathcal{B}_j} |\mathrm{II}_{\Sigma_j}|(p) = \infty$$

and so that every connected component of $\Sigma_j \cap B_{\delta}(\mathcal{B}_{\infty})$ intersects $B_{\varepsilon_j}(\mathcal{B}_{\infty})$. That we can find ε_j satisfying final condition is easily justified by combining the smooth convergence away from \mathcal{B}_{∞} to $\{x^n = 0\}$ with the varifold convergence.

Consider Σ'_j a connected component of $\Sigma_j \cap B_{\delta}(p_{\infty})$ for some $p_{\infty} \in \mathcal{B}_{\infty}$. We claim that for j sufficiently large,

(7.2)
$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,p_\infty)| < \frac{1}{4}$$

for $x \in \Sigma'_j \cap (B_{\delta}(p_{\infty}) \setminus B_{\varepsilon_j}(p_{\infty}))$. Suppose that we have proven (7.2) for each component. By the monotonicity formula and the uniform volume bound in (\beth), there must be a bounded number of such components. Thus, by taking j sufficiently large, we have that

$$|\mathrm{II}_{\Sigma_j}|(x)d_{g_j}(x,p_\infty) < \frac{1}{4}$$

for $x \in \Sigma_j \cap (B_{\delta}(p_{\infty}) \setminus B_{\varepsilon_j}(p_{\infty}))$. The inductive step (it is not hard to see that it is applicable to each connected component of $\Sigma_j \cap B_{\varepsilon_j}(p_{\infty})$, by how we chose ε_j), along with Lemma 3.1 and these bounds easily show that after passing to a subsequence each hypersurface $\Sigma_j \cap B_{\delta}(p_{\infty})$ is diffeomorphic. Passing to a further subsequence, we may arrange that each hypersurface $\Sigma_j \cap B_{\delta}(\mathcal{B}_{\infty})$ is diffeomorphic. Now, since $\Sigma_j \setminus B_{\delta/2}(\mathcal{B}_{\infty})$ converges smoothly (with finite multiplicity) to $\{x^n = 0\} \setminus B_{\delta/2}(\mathcal{B}_{\infty})$, there are only a finite number of ways that the hypersurfaces $\Sigma_j \cap B_{\delta}(\mathcal{B}_{\infty})$ could join up with $\Sigma_j \cap (B_1(0) \setminus B_{\delta}(\mathcal{B}_{\infty}))$, which is diffeomorphic to a disjoint union of finitely many copies of the "planar region" $\{x^n = 0\} \cap (B_1(0) \setminus B_{\delta/2}(\mathcal{B}_{\infty}))$. Hence, as usual it remains to prove (7.2) for each connected component Σ'_j .

The argument is similar to the one point of concentration above. If (7.2) failed, then we could choose $\delta_j \geq \varepsilon_j$ to be the smallest number so that (7.2) held for $x \in$ $\Sigma'_j \cap (B_{\delta}(p_{\infty}) \setminus B_{\delta_j}(p_{\infty}))$. The surface

$$\hat{\Sigma}'_j := \delta_j^{-1} (\Sigma'_j - p_\infty)$$

converges after passing to a subsequence to $\hat{\Sigma}'_{\infty}$. Now, we may argue exactly as in the one point case to choose $\gamma \geq 1$ so that each component of $\hat{\Sigma}'_{\infty}$ intersects $\partial B_{\gamma}(0)$ transversely in spheres. Lemma 3.1 implies that $\Sigma'_{j} \cap (B_{\delta}(p_{\infty}) \setminus B_{\gamma\delta_{j}}(p_{\infty}))$ is the union of annular regions. This implies that $\hat{\Sigma}'_{\infty}$ is connected, and is thus a plane through the origin. This contradicts the choice of δ_{j} by the same argument as before. This completes the proof in the case that $|\mathcal{B}_{\infty}| \geq 2$.

Finally, in the case that $|\mathcal{B}_{\infty}| = 1$ and $|\mathcal{B}_j| \geq 2$, we can rescale by the distance between the furthest two points of concentration. The proof proceeds just as in Proposition 4.3, as long as we prove the crucial curvature estimates from the large to small scale, as we have done above. We omit the details.

Now, to finish the proof of Theorem 1.2, we first observe that it is not restrictive to assume that the hypersurfaces are connected (the volume bounds and monotonicity formula imply that there can be at most a bounded number of connected components).

28

If Σ_j was a sequence of pairwise non-diffeomorphic connected, embedded, minimal hypersurfaces in \mathbb{R}^n with $\operatorname{vol}(\Sigma \cap B_R(0)) \leq \Lambda R^{1-n}$ and $\operatorname{index}(\Sigma) \leq I$, then because such surfaces are "regular at infinity," we may rescale and rotate the Σ_j so that outside of $B_{\tau_0/2}(0)$, the Σ_j are graphical over $\{x^n = 0\}$. This guarantees that in particular the $\Sigma_j \cap B_1(0)$ are pairwise non-diffeomorphic as well. It is not hard to show that $\Sigma_j \cap B_{r_i}(0)$ satisfies (\beth) , so the proof follows from the previous proposition.

The proof of Theorem 1.1 also follows easily from the above proposition: for $\Sigma_j \subset (M^n, g)$ as in the statement of Theorem 1.1, pairwise non-diffeomorphic, their curvature cannot be bounded. Combining the previous proposition with the usual Morse theory argument, we see that after passing to a subsequence, the Σ_j are all diffeomorphic in small fixed balls containing the points of curvature blow-up. The other portion of Σ_j converges smoothly, and there are only finitely many ways to connect the regions of large curvature to the regions of bounded curvature.

APPENDIX A. THE GENUS OF A SURFACE

Definition A.1. For Σ a non-orientable closed surface, we define the (non-orientable) genus of Σ to be

$$\operatorname{genus}(\Sigma) = \frac{1}{2}\operatorname{genus}(\widehat{\Sigma})$$

where $\widehat{\Sigma}$ is the oriented double cover.

Definition A.2. For a compact surface Σ with boundary $\partial \Sigma$ consisting of one or more closed curves, we define genus(Σ) to be the genus of the closed surface formed by gluing disks to each boundary component.

Suppose that Σ_1, Σ_2 are two oriented surfaces with boundary. If we form an oriented surface Σ by gluing together b boundary components, then from the well known formula $\chi(\Sigma) = \chi(\Sigma_1) + \chi(\Sigma_2)$, we find that

$$\operatorname{genus}(\Sigma) = \operatorname{genus}(\Sigma_1) + \operatorname{genus}(\Sigma_2) + b - 1.$$

The reader should keep in mind the example of a torus thought of as a sphere with two disks removed, glued to an annulus (along the two boundary components); neither component has any genus in the sense of Definition A.2, but obviously the torus is a genus one surface.

As a consequence of this, we find

Lemma A.3. Suppose that Σ is a properly embedded surface in $B_2(0) \subset \mathbb{R}^3$ so that there is a finite set of points $\mathcal{B} \subset B_{1/2}(0)$ which are 3ε -separated for some $\varepsilon \in (0, 1/4)$ having the following properties:

- (1) The surface Σ intersects $\partial B_{\varepsilon}(\mathcal{B})$ and $\partial B_1(0)$ transversely.
- (2) The surface $\Sigma \setminus B_{\varepsilon}(\mathcal{B})$ is topologically the union of finitely many components, each of which is topologically a disk with finitely many holes removed.
- (3) The surface $\Sigma \cap B_{\varepsilon}(\mathcal{B})$ is two-sided.
- (4) For each $p \in \mathcal{B}$, we have an upper bound r(p) on the genus of $\Sigma \cap B_{\varepsilon}(p)$ and an upper bound m(p) on the number of boundary circles $\Sigma \cap \partial B_{\varepsilon}(p)$.

Then, the genus of Σ is bounded by

$$\operatorname{genus}(\Sigma) \le \sum_{p \in \mathcal{B}} (r(p) + m(p) - 1)$$

and the number of boundary circles of $\Sigma \cap B_1(0)$ is bounded by

$$|\pi_0(\Sigma \cap \partial B_1(0))| \le \sum_{p \in \mathcal{B}} m(p).$$

Appendix B. Finite index surfaces in \mathbb{R}^3

The following theorem is a consequence of results due to Osserman [Oss64] and Fischer-Colbrie [FC85] in the two-sided case. The one sided case is due to Ros [Ros06].

Theorem B.1. Suppose that $\Sigma \hookrightarrow \mathbb{R}^3$ is a complete minimal injective immersion in \mathbb{R}^3 with finite index. Then Σ is two-sided, has finite total curvature, and is properly embedded.

Proof. By [FC85, Theorem 2] and [Ros06, Theorem 17], finite index is equivalent to finite total curvature for a complete minimal immersion in \mathbb{R}^3 . Using [Oss64], we have that Σ is conformally diffeomorphic to a punctured Riemann surface. Hence, so is the orientable double cover—this shows the orientable double cover has finite total curvature. By [Sch83c], we find that Σ is a proper embedding and is thus two-sided. \Box

This, along with the half-space theorem for minimal surfaces of finite total curvature (which is a trivial consequence of [Sch83c]) implies.

Corollary B.2. Suppose that Λ is a smooth lamination of \mathbb{R}^3 with finite index. Then, it is either a single properly embedded surface of finite total curvature or else it consists only of parallel planes, i.e. after a rotation $\Lambda = \mathbb{R}^2 \times K$ for $K \subset \mathbb{R}$ closed.

Appendix C. Removable singularity results

The following result is well known, but we indicate the proof for completeness.

Proposition C.1 (Properly embedded surfaces with curvature bounds). Suppose that (M,g) is a complete Riemannian three-manifold and $p \in M$. Suppose that for $\varepsilon > 0$, $\Sigma \subset B_{\varepsilon}(p) \setminus \{p\}$ is a properly embedded minimal surface with

$$|\mathrm{II}_{\Sigma}|(x)d_g(x,p) \le C.$$

Then, Σ smoothly extends across p, i.e. there is $\widetilde{\Sigma} \subset B_{\varepsilon}(p)$ with $\Sigma = \widetilde{\Sigma} \setminus \{p\}$.

Proof. Because Σ is proper it has finite area in $B_{\varepsilon}(p) \setminus \{p\}$. Hence, the monotonicity formula is applicable and we may consider a tangent cone to Σ at p (the tangent cone may not be unique). The assumed curvature estimate imples that any tangent cone is smooth away from $\{0\}$, so it is a single plane (possibly with multiplicity) through the origin, and blow-ups of Σ converge smoothly away from 0 to any such tangent cone. Combined with a blow-up argument, this shows that there is $\delta \in (0, \varepsilon)$ so that

$$|\mathrm{II}_{\Sigma}|(x)d_g(x,p) < \frac{1}{4}$$

for all $x \in \Sigma \cap B_{\delta}(p)$. A Morse theory argument analogous to Lemma 3.1 implies that $\Sigma \cap B_{\delta}(p)$ is the union of topological planes and annuli. Hence, it has finite Euler characteristic. A properly embedded minimal surface with finite Euler characteristic is well known to extend across a point singularity, cf. [CS85, Proposition 1].

We will make use of the following Bernstein-type result due to Gulliver–Lawson [GL86]; see also [MPR13, Lemma 3.3] and [CM15, Lemma A.26].

Theorem C.2 (Gulliver–Lawson's Bernstein theorem). Suppose that $\varphi : \Sigma \to \mathbb{R}^3 \setminus \{0\}$ is a non-empty two-sided stable minimal immersion which is complete away from $\{0\}$. Then the trace of φ is a plane.

Using this, we show the following removable singularity result for two-sided stable laminations. The fundamental strategy is somewhat similar to [MPR13], but thanks to the stability hypothesis (which is considerably stronger than the assumptions in [MPR13]), we are able to give a relatively short argument, inspired by ideas in [CCE15].

Proposition C.3 (Removable singularities for two-sided stable laminations). Suppose that (M,g) is a complete Riemannian three-manifold and $p \in M$. Suppose that for some $\varepsilon > 0$, $\mathcal{L} \subset B_{\varepsilon}(p) \setminus \{p\}$ is a minimal lamination with the property that any leaf $L \subset \mathcal{L}$ has stable universal cover. Then, there is a smooth lamination $\widetilde{\mathcal{L}} \subset B_{\varepsilon}(p)$ so that $\mathcal{L} = \widetilde{\mathcal{L}} \setminus \{p\}$.

Proof. Because the claim is purely local, at several points we will replace \mathcal{L} with its intersection with some smaller ball $B_{\varepsilon'}(p)$. For simplicity, we will not relabel the resulting immersion or ball. Furthermore, we will always work in a normal coordinate system around p (where we can assume the metric to be sufficiently close to Euclidean, by taking $\varepsilon > 0$ sufficiently small).

Observe that, taking ε smaller if necessary, by Schoen's curvature estimates [Sch83a], there is C > 0 so that $|\Pi_{\mathcal{L}}|(x)d_g(x,p) \leq C$ for all $x \in \mathcal{L}$. Hence, for any $\rho_j \to \infty$, passing to a subsequence, the laminations $\rho_j(\mathcal{L}_j - p)$ converge to a smooth lamination \mathcal{L}_{∞} of $\mathbb{R}^3 \setminus \{0\}$ away from $\{0\}$. Moreover each leaf in \mathcal{L}_{∞} has stable universal cover and is complete away from $\{0\}$. Hence by Theorem C.2, after rotating, $\mathcal{L}_{\infty} = (\mathbb{R}^2 \times K) \setminus \{0\}$ for some closed set $K \subset \mathbb{R}$.

Thus, taking $\varepsilon > 0$ sufficiently small, we may guarantee that

(C.1)
$$|\mathrm{II}_{\mathcal{L}}|(x)d_g(x,p) < \frac{1}{4}$$

for all $x \in \mathcal{L}$.

Now, pick any leaf $L \subset \mathcal{L}$ so that p is in the (set-theoretic) closure of L. We claim that for any $\rho_j \to 0$, passing to a subsequence and rotating the coordinate chart, the surfaces $L_j := \rho_j L$ converge to the lamination $(\mathbb{R}^2 \times \{0\}) \setminus \{0\} \subset \mathbb{R}^3 \setminus \{0\}$. To prove this, by our above argument, it is sufficient to show that there is not another plane $\Pi = \mathbb{R}^2 \times \{z\}$ in the lamination limit of $\rho_j L$. If such a plane did exist, then by curvature estimates, we would have locally smooth convergence to Π (because $0 \notin \Pi$). At the scale of L, this would imply that there was some $\delta \in (0, \varepsilon)$ sufficiently small, so that $L \cap B_{\delta}(p) \setminus \{p\}$ contains a properly embedded component D diffeomorphic to a disk, intersecting $\partial B_{\delta}(p)$ transversely. Now, by the curvature estimates (C.1), we claim that a Morse theory argument along the lines of Lemma 3.1 shows that $L \setminus B_{\delta}(p)$ contains an annular region connecting the disk D to $\partial B_{\varepsilon}(0)$. Given this, because L is connected (by definition) we see that it must be the union of D with this annular region. This implies that p cannot be in the closure of L, a contradiction. We emphasize that L is not assumed to be proper, so we cannot simply apply Lemma 3.1. However, the given curvature estimates and resulting compactness properties of the leaves are sufficient to handle the lack of properness in the proof of the mountain pass lemma; this has been carried out in detail in [CCE15, Appendix E] in a completely analogous situation.

Thus, we may take $\delta \in (0, \varepsilon/3)$ sufficiently small so that after rotating the normal coordinate system, $L \cap (B_{3\delta}(p) \setminus B_{\delta}(p))$ intersects $\partial B_{2\delta}(p)$ transversely, is contained in a $\delta/10$ neighborhood of the coordinate plane $\mathbb{R}^2 \times \{0\}$, and is a multigraph over this plane (cf. [CCE15, Lemma 4.1]). Hence, the intersection $L \cap \partial B_{2\delta}(p)$ is the union of simple closed curves and injectively immersed curves which "spiral near the equator" of the sphere $\partial B_{2\delta}(p)$. First, assume that $L \cap \partial B_{2\delta}(p)$ contains a simple closed curve. Then, by the curvature estimate (C.1) and the Morse theory argument used above, we see that $L \cap \partial B_{3\delta}(p)$ is a properly embedded annulus in $B_{3\delta}(p) \setminus \{p\}$. Hence, L extends across $\{p\}$ by Proposition C.1.

Thus, it remains to consider the case that $L \cap \partial B_{2\delta}(p)$ consists of one or more spiraling curves. Our argument here is analogous to the technique of passing to the top sheet in [CCE15, Proposition 4.2]. We have seen that $L \cap \partial B_{2\delta}(p)$ is contained in an $\delta/10$ neighborhood of the equator $(\mathbb{R}^2 \times \{0\} \cap \partial B_{2\delta}(p))$. Taking a sequence of points $w_j \in L \cap \partial B_{2\delta}(p)$ with $x^3(w_j)$ approaching $\sup_{w \in L \cap B_{2\delta}(p)} x^3(w)$, after passing to a subsequence, the points w_j converge to w', which lies in a "top sheet" $L' \subset \mathcal{L}$. By construction, $L' \cap \partial B_{2\delta}(p)$ will contain a simple closed curve, and thus the Morse theory argument guarantees that it $L' \cap B_{3\delta}(p)$ is a properly embedded topological disk or annulus and thus, by Proposition C.1, L' extends across $\{p\}$. Similarly, we may pass to the bottom sheet to find a properly embedded $L'' \subset \mathcal{L}$ which extends across $\{p\}$. Note that by construction $L' \neq L''$ (otherwise, there could not be any spiraling).

Note that by the maximum principle, it cannot happen that both L' and L'' contain p (because they are smoothly, properly embedded in $B_{3\delta}(p)$), so we assume that $p \notin L'$. However, this leads to a contradiction as follows: by construction and the curvature estimates, we can find a sufficiently small tubular neighborhood \mathcal{U} of $L' \cap B_{3\delta}(p)$ so that $L \cap \mathcal{U}$ is a multigraph over $L' \cap B_{3\delta}(p)$. Because $L' \cap B_{3\delta}(p)$ is a disk (and thus is simply connected), this shows that at least one component of $L \cap \mathcal{U}$ must be a disk, contradicting the assumed spiraling behavior of L.

Remark C.4. We remark that by combining the curvature estimates [Sch83a, Ros06] with the removable singularity result [MPR13], the following strengthened version of Proposition C.3 holds: suppose that $\mathcal{L} \subset B_{\varepsilon}(p) \setminus \{p\}$ is a lamination so that every leaf has a cover that is stable. Then, \mathcal{L} extends smoothly across $\{p\}$. Note that that this version is compatible with one-sided stability of leaves, while Proposition C.6 requires two-sided stability.¹⁴

¹⁴We emphasize that one-sided stability does not necessarily imply that the universal cover is stable; consider $\mathbb{R}P^2$ in $\mathbb{R}P^3$.

We have not been able to find a self contained proof of this strengthened assertion, due to the fact that we do not know if the one-sided version of Theorem C.2 holds. It would thus be interesting to know if there can be a non-flat, one-sided stable immersion in \mathbb{R}^3 that is complete away from the origin. Note that the surface cannot be injectively immersed, because then by the curvature estimates [Ros06], one could take the lamination closure away from {0} and then apply [MPR13].

Fortunately, for our purposes, the removable singularity result contained in Proposition C.3 is sufficient, as we prove that two-sidedness "propagates outwards" in the event of index concentration (cf., Proposition 4.2).

Corollary C.5. Suppose that $\mathcal{B} \subset \mathbb{R}^3$ is a finite set of points and $\Lambda \subset \mathbb{R}^3 \setminus \mathcal{B}$ is a smooth lamination of $\mathbb{R}^3 \setminus \mathcal{B}$ so that

$$|\mathrm{II}_{\Lambda}|(x)d_{\mathbb{R}^3}(x,\mathcal{B}) \leq C.$$

Then, either Λ extends smoothly across the points \mathcal{B} or it contains a plane.

Proof. If each leaf is properly embedded in $\mathbb{R}^3 \setminus \mathcal{B}$, then Proposition C.1 implies that Λ extends across \mathcal{B} . Otherwise, Λ contains some limit leaf L, which has stable universal cover by standard arguments (cf. [MR06, Lemma A.1]). Then the closure of L in Λ is a non-empty lamination consisting entirely of leaves with stable universal cover. Proposition C.3 then guarantees that the closure of L extends smoothly across \mathcal{B} . Hence, we see that L must be a plane, by [FCS80, dCP79, Pog81].

Finally, we need the following removable singularity result valid in higher dimensions.

Proposition C.6. Suppose that (M^n, g) is a complete Riemannian n-dimensional manifold, for $4 \le n \le 7$ and for some $\varepsilon > 0$, $\Sigma \subset B_{\varepsilon}(p) \setminus \{p\}$ is a properly embedded stable minimal hypersurface. Then, Σ smoothly extends across p.

Proof. Stability of the hypersurface implies that $|II_{\Sigma}|(x)d_g(x,p) \leq C$. This follows from a blow-up argument as in Lemma 2.4, based on [SSY75, SS81] and the fact that a properly embedded hypersurface in \mathbb{R}^n is two-sided (cf. [Sam69]).

As usual, the claim is purely local, so there is no harm with assuming that $\varepsilon > 0$ is sufficiently small. The curvature estimates and properness guarantee that $\operatorname{vol}_g(\Sigma) < \infty$. Thus, the monotonicity formula allows us to consider the tangent cones to (the varifold closure of) Σ at p. The curvature estimates guarantee all of the tangent cones have smooth, compact, connected, cross section in \mathbb{S}^{n-1} . It is well known that (by Alexander duality, cf. [Sam69]) compact embedded hypersurfaces separate \mathbb{S}^{n-1} , and are thus twosided. Thus, the tangent cones themselves are two-sided. Because the rescalings of Σ_j converge smoothly away from the origin to the tangent cones, we may see that the cones are stable precisely in the sense needed to apply [Sim68]. This allows us to conclude that all tangent cones to Σ at p are hyperplanes (possibly with multiplicity).

In particular, taking $\varepsilon > 0$ smaller if necessary, we may arrange that

$$|\mathrm{II}_{\Sigma}|(x)d_g(x,p) \le \frac{1}{4}.$$

Then, using a Morse theoretic argument along the lines of Lemma 2.4 (and taking $\varepsilon > 0$ smaller if necessary), we may arrange that each of the (bounded number of) components of Σ is diffeomorphic to $\mathbb{S}^{n-2} \times (0, 1)$ and each component intersects $\partial B_s(p)$ transversely

in a connected submanifold for $s \leq \varepsilon$. From this, it is not hard to see that any tangent cone to (the varifold closure of) the hypersurface $\tilde{\Sigma}$ at p is a multiplicity one plane. Thus, by Allard's theorem [All72], $\tilde{\Sigma}$ extends smoothly across p. Using the maximum principle, we thus see that there can be only one component of Σ whose closure includes p. This completes the proof.

APPENDIX D. EXAMPLES OF DEGENERATION

We give examples to illustrate the "one point of concentration" and "multiple points of concentration" discussed in Propositions 4.2 and 4.3. Recall that (see [Cos84, HM85, HK97]) the Costa surface $\Sigma^{(1)} \subset \mathbb{R}^3$ is an embedded minimal surface with genus one and three ends (one flat end, and two catenoidal ends) Furthermore, Hoffman–Meeks have shown that it is possible to deform the flat end of Σ_1 into a catenoidal end, producing a family $\Sigma^{(t)}$ for $t \geq 1$ of embedded genus three embedded minimal surfaces with three catenoidal ends. As $t \to \infty$, the logarithmic growth of middle catenoidal end approaches that of the other end pointing in the same direction. See [HK97, Figure 3.2] or Figure 1 for an illustration of the deformation family for t large.

The exact index of $\Sigma^{(t)}$ seems to be unknown for t > 1 (note that $index(\Sigma^{(1)}) = 5$ by [Nay93]). However, because the family $\Sigma^{(t)}$ has uniformly bounded total curvature, the main result of [Tys87] implies that $index(\Sigma^{(t)}) \leq I$ for some $I \in \mathbb{N}$. We will always assume that Σ^t is scaled so that the second fundamental form's maximal norm is equal to 1 and so that the line $\{x^1 = x^2 = 0\}$ is the intersection of the two planes of reflection symmetry.



FIGURE 1. This depicts the Hoffman–Meeks deformation $\Sigma^{(t)}$ of the Costa surface for t large. The surface looks like three planes joined by three catenoidal necks. The marked points are the points of curvature concentration, and at the scale of curvature around any of them, $\Sigma^{(t)}$ is geometrically close to a catenoid. By choosing the scaling appropriately, one may arrange that the point of concentration stay at a bounded distance, or all converge to the origin.

First, to illustrate the case of one point of concentration simply consider

$$\Sigma_j = \left(\frac{1}{j}\Sigma^{(1)}\right) \cap B_{r_j}(0)$$

for some $r_j \to \infty$. As $r \to \infty$, this converges smoothly away from the origin to the plane $\mathbb{R}^2 \times \{0\}$ with multiplicity 3. Clearly this satisfies (\aleph) with one point of curvature concentration p_j converging to 0 as $j \to \infty$. Rescaling the sequence at the scale of curvature around p_j simply finds Σ_1 .

Second, to illustrate the various possibilities for multiple points of concentration, we must describe the behavior of $\Sigma^{(t)}$ as $t \to \infty$ more precisely. One may show that for $\rho_i \to 0$ sufficiently quickly,

$$\Sigma_j := \left(\rho_j \Sigma^{(j)}\right) \cap B_{r_j}(0)$$

looks like three nearby disks, with the middle disk jointed to the bottom disk by two catenoidal necks in equal and opposite directions from the origin and the middle disk joined to the top disk by a single catenoidal neck near the origin. This is well illustrated in [HK97, Figure 3.2]; see also Figure 1. To establish this picture rigorously, one may appeal to [Ros95, Theorem 2] and the fact that the catenoid is the only nonflat embedded minimal surface $\hat{\Sigma} \subset \mathbb{R}^3$ with total curvature $\int_{\hat{\Sigma}} \kappa > -12\pi$ (cf. [HK97, Theorem 3.1]).

In particular, the blow-up set \mathcal{B}_j has $|\mathcal{B}_j| = 3$ and rescaling around any such point is a catenoid. However, if we chose $\rho_j \to 0$ sufficiently quickly, \mathcal{B}_j converges to $\mathcal{B}_{\infty} = \{0\}$ as $j \to \infty$. On the other hand, if $\rho_j \to 0$ at precisely the correct rate, it is clear that (after a rotation) \mathcal{B}_j converges to $\{(0,0,0), (\pm 1,0,0)\} \subset \mathbb{R}^3$. This example, and considerably more refined examples are discussed in great detail in [Tra02].

References

- [All72] William K. Allard, On the first variation of a varifold, Ann. of Math. (2) 95 (1972), 417–491.
 MR 0307015 (46 #6136)
- [BS15] Reto Buzano and Ben Sharp, Qualitative and quantitative estimates for minimal hypersurfaces with bounded index and area, preprint, available at http://arxiv.org/abs/1512.01047 (2015).
- [BW69] Glen E. Bredon and John W. Wood, Non-orientable surfaces in orientable 3-manifolds, Invent. Math. 7 (1969), 83–110. MR 0246312 (39 #7616)
- [Car15] Alessandro Carlotto, Generic finiteness of minimal surfaces with bounded Morse index, preprint, available at http://arxiv.org/abs/1509.07101 (2015).
- [CCE15] Alessandro Carlotto, Otis Chodosh, and Michael Eichmair, Effective versions of the positive mass theorem, preprint, available at http://arxiv.org/abs/1503.05910 (2015).
- [CDL05] Tobias H. Colding and Camillo De Lellis, Singular limit laminations, Morse index, and positive scalar curvature, Topology 44 (2005), no. 1, 25–45. MR 2103999 (2005h:53102)
- [CG14] Tobias H. Colding and David Gabai, Effective finiteness of irreducible Heegaard splittings of non Haken 3-manifolds, preprint, available at http://arxiv.org/abs/1411.2509 (2014).
- [CH06] Tobias H. Colding and Nancy Hingston, Geodesic laminations with closed ends on surfaces and Morse index; Kupka-Smale metrics, Comment. Math. Helv. 81 (2006), no. 3, 495–522. MR 2250851 (2007f:53037)
- [CM00] Tobias H. Colding and William P. Minicozzi, II, Embedded minimal surfaces without area bounds in 3-manifolds, Geometry and topology: Aarhus (1998), Contemp. Math., vol. 258, Amer. Math. Soc., Providence, RI, 2000, pp. 107–120. MR 1778099 (2001i:53012)
- [CM14] Otis Chodosh and Davi Maximo, On the topology and index of minimal surfaces, preprint, available at http://arxiv.org/abs/1405.7356 (2014).
- [CM15] Tobias H. Colding and William P. Minicozzi, II, The space of embedded minimal surfaces of fixed genus in a 3-manifold V; fixed genus, Ann. of Math. (2) 181 (2015), no. 1, 1–153. MR 3272923

- [Cos84] Celso J. Costa, Example of a complete minimal immersion in R³ of genus one and three embedded ends, Bol. Soc. Brasil. Mat. 15 (1984), no. 1-2, 47–54. MR 794728 (87c:53111)
- [CS85] Hyeong In Choi and Richard Schoen, The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature, Invent. Math. 81 (1985), no. 3, 387–394. MR 807063 (87a:58040)
- [CW83] Hyeong In Choi and Ai Nung Wang, A first eigenvalue estimate for minimal hypersurfaces,
 J. Differential Geom. 18 (1983), no. 3, 559–562. MR 723817 (85d:53028)
- [dCP79] Manfredo do Carmo and Chiakuei Peng, Stable complete minimal surfaces in \mathbb{R}^3 are planes, Bull. Amer. Math. Soc. (N.S.) **1** (1979), no. 6, 903–906. MR 546314 (80j:53012)
- [Dea03] Brian Dean, Compact embedded minimal surfaces of positive genus without area bounds, Geom. Dedicata 102 (2003), 45–52. MR 2026836 (2005d:53011)
- [EM08] Norio Ejiri and Mario Micallef, Comparison between second variation of area and second variation of energy of a minimal surface, Adv. Calc. Var. 1 (2008), no. 3, 223–239. MR 2458236 (2009j:58019)
- [FC85] Doris Fischer-Colbrie, On complete minimal surfaces with finite Morse index in threemanifolds, Invent. Math. 82 (1985), no. 1, 121–132. MR 808112 (87b:53090)
- [FCS80] Doris Fischer-Colbrie and Richard Schoen, The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature, Comm. Pure Appl. Math. 33 (1980), no. 2, 199–211. MR 562550 (81i:53044)
- [FHS83] Michael Freedman, Joel Hass, and Peter Scott, Least area incompressible surfaces in 3manifolds, Invent. Math. 71 (1983), no. 3, 609–642. MR 695910 (85e:57012)
- [GL86] Robert Gulliver and H. Blaine Lawson, Jr., The structure of stable minimal hypersurfaces near a singularity, Geometric measure theory and the calculus of variations (Arcata, Calif., 1984), Proc. Sympos. Pure Math., vol. 44, Amer. Math. Soc., Providence, RI, 1986, pp. 213– 237. MR 840275 (87g:53091)
- [HI01] Gerhard Huisken and Tom Ilmanen, The inverse mean curvature flow and the Riemannian Penrose inequality, J. Differential Geom. 59 (2001), no. 3, 353–437. MR 1916951 (2003h:53091)
- [HK97] David Hoffman and Hermann Karcher, Complete embedded minimal surfaces of finite total curvature, Geometry, V, Encyclopaedia Math. Sci., vol. 90, Springer, Berlin, 1997, pp. 5–93. MR 1490038 (98m:53012)
- [HM85] D. Hoffman and W. H. Meeks, III, A complete embedded minimal surface in R³ with genus one and three ends, J. Differential Geom. 21 (1985), no. 1, 109–127. MR 806705 (87d:53008)
- [Jac70] William Jaco, Surfaces embedded in $M^2 \times S^1$, Canad. J. Math. **22** (1970), 553–568. MR 0267596 (42 #2498)
- [Kra09] Joel Kramer, *Embedded minimal spheres in 3-manifolds*, Ph.D. thesis, The Johns Hopkins University, 2009.
- [LZ15] Haozhao Li and Xin Zhou, Existence of minimal surfaces of arbitrary large morse index, preprint, available at http://arxiv.org/abs/1504.00970 (2015).
- [MN13] Fernando C. Marques and André Neves, Existence of infinitely many minimal hypersurfaces in positive Ricci curvature, preprint, available at http://arxiv.org/abs/1311.6501 (2013).
- [MPR13] William H. Meeks, III, Joaquin Perez, and Antonio Ros, *Local removable singularity theorems* for minimal laminations, preprint, available at http://arxiv.org/abs/1308.6439 (2013).
- [MR05] William H. Meeks, III and Harold Rosenberg, *The uniqueness of the helicoid*, Ann. of Math.
 (2) 161 (2005), no. 2, 727–758. MR 2153399 (2006f:53012)
- [MR06] _____, The minimal lamination closure theorem, Duke Math. J. **133** (2006), no. 3, 467–497. MR 2228460 (2007e:53007)
- [Nay93] Shin Nayatani, Morse index and Gauss maps of complete minimal surfaces in Euclidean 3-space, Comment. Math. Helv. 68 (1993), no. 4, 511–537. MR 1241471 (95b:58039)
- [Oss64] Robert Osserman, Global properties of minimal surfaces in E^3 and E^n , Ann. of Math. (2) 80 (1964), 340–364. MR 0179701 (31 #3946)
- [Pit76] Jon T. Pitts, Existence and regularity of minimal surfaces on Riemannian manifolds, Bull. Amer. Math. Soc. 82 (1976), no. 3, 503–504.

- [Pog81] Aleksei V. Pogorelov, On the stability of minimal surfaces, Dokl. Akad. Nauk SSSR 260 (1981), no. 2, 293–295. MR 630142 (83b:49043)
- [Ros95] Antonio Ros, Compactness of spaces of properly embedded minimal surfaces with finite total curvature, Indiana Univ. Math. J. 44 (1995), no. 1, 139–152. MR 1336435 (96k:53011)
- [Ros06] _____, One-sided complete stable minimal surfaces, J. Differential Geom. 74 (2006), no. 1, 69–92. MR 2260928 (2007g:53008)
- [Sam69] Hans Samelson, Orientability of hypersurfaces in \mathbb{R}^n , Proc. Amer. Math. Soc. **22** (1969), 301–302. MR 0245026 (39 #6339)
- [Sch83a] Richard Schoen, Estimates for stable minimal surfaces in three-dimensional manifolds, Seminar on minimal submanifolds, Ann. of Math. Stud., vol. 103, Princeton Univ. Press, Princeton, NJ, 1983, pp. 111–126. MR 795231 (86j:53094)
- [Sch83b] Richard M. Schoen, Uniqueness, symmetry, and embeddedness of minimal surfaces, J. Differential Geom. 18 (1983), no. 4, 791–809 (1984). MR 730928 (85f:53011)
- [Sch83c] _____, Uniqueness, symmetry, and embeddedness of minimal surfaces, J. Differential Geom.
 18 (1983), no. 4, 791–809 (1984). MR 730928 (85f:53011)
- [Sha15] Ben Sharp, Compactness of minimal hypersurfaces with bounded index, preprint, available at http://arxiv.org/abs/1501.02703 (2015).
- [Sim68] James Simons, Minimal varieties in riemannian manifolds, Ann. of Math. (2) 88 (1968), 62–105. MR 0233295 (38 #1617)
- [SS81] Richard Schoen and Leon Simon, Regularity of stable minimal hypersurfaces, Comm. Pure Appl. Math. 34 (1981), no. 6, 741–797. MR 634285 (82k:49054)
- [SSY75] Richard Schoen, Leon Simon, and Shing-Tung Yau, Curvature estimates for minimal hypersurfaces, Acta Math. 134 (1975), no. 3-4, 275–288. MR 0423263 (54 #11243)
- [SY79] R. Schoen and Shing Tung Yau, Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature, Ann. of Math. (2) 110 (1979), no. 1, 127–142. MR 541332 (81k:58029)
- [SY82] Richard Schoen and Shing Tung Yau, Complete three-dimensional manifolds with positive Ricci curvature and scalar curvature, Seminar on Differential Geometry, Ann. of Math. Stud., vol. 102, Princeton Univ. Press, Princeton, N.J., 1982, pp. 209–228. MR 645740 (83k:53060)
- [SY83] Richard Schoen and S. T. Yau, The existence of a black hole due to condensation of matter, Comm. Math. Phys. 90 (1983), no. 4, 575–579. MR 719436 (84k:83005)
- [Tra02] Martin Traizet, An embedded minimal surface with no symmetries, J. Differential Geom. 60 (2002), no. 1, 103–153. MR 1924593 (2004c:53008)
- [Tra04] _____, A balancing condition for weak limits of families of minimal surfaces, Comment. Math. Helv. 79 (2004), no. 4, 798–825. MR 2099123 (2005g:53017)
- [Tys87] Johan Tysk, Eigenvalue estimates with applications to minimal surfaces, Pacific J. Math. 128 (1987), no. 2, 361–366. MR 888524 (88i:53102)
- [Tys89] _____, Finiteness of index and total scalar curvature for minimal hypersurfaces, Proc. Amer. Math. Soc. 105 (1989), no. 2, 429–435. MR 946639 (89g;53093)
- [Whi87] B. White, Curvature estimates and compactness theorems in 3-manifolds for surfaces that are stationary for parametric elliptic functionals, Invent. Math. 88 (1987), no. 2, 243–256. MR 880951 (88g:58037)
- [Whi91] Brian White, The space of minimal submanifolds for varying Riemannian metrics, Indiana Univ. Math. J. 40 (1991), no. 1, 161–200. MR 1101226 (92i:58028)
- [Whi15] Brian White, On the bumpy metrics theorem for minimal submanifolds, preprint, available at http://arxiv.org/abs/1503.01803 (2015).
- [YY80] Paul C. Yang and Shing Tung Yau, Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 7 (1980), no. 1, 55–63. MR 577325 (81m:58084)

OTIS CHODOSH, DANIEL KETOVER, AND DAVI MAXIMO

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, CAMBRIDGE UNIVERSITY *E-mail address:* oc249@cam.ac.uk

 $\label{eq:constraint} \begin{array}{l} \text{Department of Mathematics, Princeton University} \\ \textit{E-mail address: dketover@math.princeton.edu} \end{array}$

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY *E-mail address*: maximo@math.stanford.edu

38