

Read Fulton, Chapter 7, sections 4-5.

1. a) In Hartshorne, Chapter IV, do problems 1.5, 1.7, 2.5.  
 b) (Optional) In Fulton, Chapter 7, do problem 7.12 in Section 7.4; and problems 7.17 and 7.21(a) in Section 7.5
2. Let  $k$  be an algebraically closed field of characteristic  $p \neq 0$ .  
 (a) Find the genus of the smooth completion  $Y$  of the affine curve  $Y^\circ$  given by  $y^p - y = x$ . [Hint: Project onto the  $y$ -axis.]  
 (b) Let  $X$  be the projective  $x$ -line over  $k$ . Show that the projection map  $Y^\circ \rightarrow \mathbb{A}_k^1$  onto the  $x$ -axis extends to a morphism  $Y \rightarrow X$ , and find where this morphism ramifies.  
 (c) Using (a) and (b), conclude that the Hurwitz formula for covers  $Y \rightarrow X$  of degree  $n$ , i.e.  $2g_Y - 2 = n(2g_X - 2) + \sum_{Q \in X} (e_Q - 1)$ , does not carry over to wildly ramified covers in characteristic  $p > 0$ .
3. Let  $k$  be a field.  
 a) Show that if  $k$  is algebraically closed and of characteristic 0, then  $\mathbb{A}_k^1$  is simply connected (i.e. it has no non-trivial connected étale covers) .  
 b) Prove the converse of (a).
4. Let  $\pi : Y \rightarrow X$  be a covering space map of Riemann surfaces (in the complex metric topology), where  $X$  is compact and of topological genus 1.  
 (a) Show that if  $\pi$  is of finite degree, then  $Y$  is also compact, and also of topological genus 1.  
 (b) Show that the above conclusion fails if  $\pi$  is of infinite degree. [Hint: Consider the universal cover.] Why doesn't this phenomenon arise in the algebraic situation (i.e. for morphisms)?
5. Let  $Y$  be a smooth connected curve over a field  $k$ . Let  $G$  be a finite group that acts faithfully on  $Y$  as a group of automorphisms. Suppose that the order of  $G$  is not divisible by the characteristic of  $k$ .  
 a) Show that there is a smooth curve  $X$  and a morphism  $\pi : Y \rightarrow X$  which is a Galois branched cover having Galois (covering) group  $G$ , such that the fibres of  $\pi$  are the orbits of  $G$ . We call  $X$  the *quotient*  $Y/G$ . [Hint: First handle the affine case  $Y = \text{Spec } S$ , by taking  $X = \text{Spec } S^G$ , where  $S^G$  is the subring of  $G$ -invariants of  $S$ .]  
 b) Suppose that  $k$  is algebraically closed, and let  $Q \in Y$ . Show that  $Q$  is stabilized by some element of  $G$  other than the identity if and only if  $Q$  is a ramification point of  $Y \rightarrow X$ . Show that the number of elements of  $G$  that stabilize  $Q$  (including the identity) is equal to  $e_Q$ .  
 c) Let  $k = \mathbb{C}$ , and let  $Y = \mathbb{P}_{\mathbb{C}}^1$ , the Riemann sphere. Let  $G$  be the symmetry group of the solid tetrahedron  $T$ , where  $T$  is viewed as inscribed in the sphere. Consider the action of  $G$  on  $Y$  that is induced by the action of  $G$  on  $T$ . Describe the resulting cover  $Y \rightarrow X = Y/G$  by giving its degree, the number of branch and ramification points, and the ramification indices; giving the genus of  $X$ ; and checking that the Hurwitz formula holds.